## ENNAKKOMATERIAALI

• Artikkeli:

McClements DJ, Öztürk B. 2021. Utilization of Nanotechnology to Improve the Handling, Storage and Biocompatibility of Bioactive Lipids in Food Applications. *Foods*, 10, 365. https://doi.org/10.3390/foods10020365

- Liite 1: sanasto
- Liite 2: korjatut rakennekaavat (kts. alla oleva lisätieto)

Ennakkomateriaalina olevan artikkelin kuvassa 1 (figure1) on osa rakenteista esitetty virheellisesti. Korjatut rakennekaavat löytyvät artikkelitiedoston lopusta liitteestä 2.





# **Utilization of Nanotechnology to Improve the Handling, Storage and Biocompatibility of Bioactive Lipids in Food Applications**

David Julian McClements <sup>1,2,\*</sup> and Bengü Öztürk <sup>3</sup>

- <sup>1</sup> Department of Food Science, University of Massachusetts Amherst, Amherst, MA 01003, USA
- <sup>2</sup> Department of Food Science & Bioengineering, Zhejiang Gongshang University, Hangzhou 310018, China
- <sup>3</sup> Department of Food Engineering, Faculty of Engineering, Yeditepe University, Istanbul 34755, Turkey
  - Correspondence: mcclements@foodsci.umass.edu

Abstract: Bioactive lipids, such as fat-soluble vitamins, omega-3 fatty acids, conjugated linoleic acids, carotenoids and phytosterols play an important role in boosting human health and wellbeing. These lipophilic substances cannot be synthesized within the human body, and so people must include them in their diet. There is increasing interest in incorporating these bioactive lipids into functional foods designed to produce certain health benefits, such as anti-inflammatory, antioxidant, anticancer and cholesterol-lowering properties. However, many of these lipids have poor compatibility with food matrices and low bioavailability because of their extremely low water solubility. Moreover, they may also chemically degrade during food storage or inside the human gut because they are exposed to certain stressors, such as high temperatures, oxygen, light, moisture, pH, and digestive/metabolic enzymes, which again reduces their bioavailability. Nanotechnology is a promising technology that can be used to overcome many of these limitations. The aim of this review is to highlight different kinds of nanoscale delivery systems that have been designed to encapsulate and protect bioactive lipids, thereby facilitating their handling, stability, food matrix compatibility, and bioavailability. These systems include nanoemulsions, solid lipid nanoparticles (SLNs), nanostructured lipid carriers (NLCs), nanoliposomes, nanogels, and nano-particle stabilized Pickering emulsions.

Keywords: nanotechnology; bioactive lipids; storage stability; bioavailability; food applications

## 1. Introduction

Bioactive lipids are oil-soluble compounds that may provide certain health benefits to humans when consumed in appropriate amounts, i.e., doses and frequencies. Depending on their mechanisms of action, they may boost the immune system, reduce inflammation, improve bone health, eye and brain functions, reduce coronary heart diseases, and act as antioxidants and anticarcinogens [1-4]. Some of the most commonly studied bioactive lipids that may promote human health and well-being are phytosterols (e.g.,  $\beta$ -sitosterol, stigmasterol, campesterol), fat soluble vitamins (e.g., vitamins A, D, E, K), carotenoids (e.g.,  $\beta$ -carotene, lycopene, lutein, zeaxanthin, astaxanthin) and essential fatty acids (e.g., ALA, EPA, DHA, CLA,) [5,6] (Figure 1). As well as their traditional bioactivities (Section 2), a number of studies have recently shown that consumption of some of these bioactive lipids may also provide protection against human coronavirus SARS-CoV-2. For instance, ingestion of EPA and DHA has been reported to reduce inflammation, cytokine storm and blood coagulation, which might play role an important role in the recovery of patients with Covid-19 infection [7–9]. Health professionals have also recommended Covid-19 patients to take vitamin D to modulate respiratory tract disorders, boost immune cell proliferation and activity, and promote pulmonary ACE2 expression mechanisms [10].



Citation: McClements, D.J.; Öztürk, B. Utilization of Nanotechnology to Improve the Handling, Storage and Biocompatibility of Bioactive Lipids in Food Applications. *Foods* **2021**, *10*, 365. https://doi.org/10.3390/ foods10020365

Academic Editors: Luis M. Rodríguez-Alcalá, Lígia Pimentel and Susana Vidigal Received: 11 January 2021 Accepted: 5 February 2021 Published: 8 February 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/).



**Figure 1.** Chemical structures of bioactive lipids: (**a**) fat soluble vitamins, carotenoids, phytosterols, (**b**) polyunsaturated fatty acids.

Most bioactive lipids are not synthesized in the human body and so they have to be obtained from food sources, such as nuts, seeds, grains, eggs, meat, fish, tomato, fruits, and vegetables [1,11–13]. However, many people do not get enough of these bioactive compounds from their regular diets due to their low dietary levels, tendency to degrade during processing or storage, and their poor stability, solubility, and absorption characteristics within the human gut [14–17]. The food industry is therefore exploring methods of fortifying foods with stable and bioavailable forms of these bioactive lipids.

Nanotechnology has emerged as a powerful means of encapsulating, protecting, and delivering bioactive substances in foods, thereby improving their efficacy [18,19]. These nanoscale delivery systems should be constructed entirely from food grade ingredients and should be designed to provide resistance against the elevated temperatures, light levels, and oxygen levels they may be exposed to during food processing and storage. They should also be designed to be stable within the specific food matrix that they are utilized within.

The aim of this review is to highlight the potential of nanoscale delivery systems for improving the handling, storage, and bioavailability of bioactive lipids in foods. Initially, the different kinds of bioactive lipids typically used in foods are introduced, and then the various types of nanoscale delivery systems that can be used to encapsulate them are reviewed. Examples of how these delivery systems are utilized to improve the handling, storage, and bioavailability of bioactive lipids are then given.

## 2. Bioactive Lipids

#### 2.1. Fat Soluble Vitamins

Fat soluble vitamins are hydrophobic micronutrients that humans must consume to maintain various body functions and ensure health and wellbeing (Figure 1a) [6,20]. Vitamin A (all-trans retinol) is found in a variety of foods, with the chemical form depending on the source. Retinol is mainly found in plant-based foods while retinyl esters are found in animal-based foods. Vitamin A is known to improve visual health, help reproduction, boost the immune system, and promote growth [21]. Vitamin D is a steroid hormone that comes in two major forms: vitamin D<sub>2</sub> (ergocalciferol) and D<sub>3</sub> (cholecalciferol). In the body, the most bioactive form of vitamin D is 1,25-dihydroxyvitamin D<sub>3</sub>, which plays an important role in modulating the immune system, calcium metabolism, and the maintenance of bone and teeth health [22]. Vitamin E is comprised of a number of tocopherols ( $\alpha$ -,  $\beta$ -,  $\gamma$ -, and  $\delta$ -tocopherol) and tocotrienols ( $\alpha$ -,  $\beta$ -,  $\gamma$ -, and  $\delta$ tocotrienol) that are essential for human health and performance. They also act as natural antioxidants and help prevent cancer, heart disease, diabetes, and skin diseases [23]. Vitamin K is a group of hydrophobic compounds (phylloquinones and menaquinones) that all have a methylated naphthoquinone ring system, which are substituted with isoprenoid and isoprene side chains, respectively. Vitamin K is a micronutrient whose ingestion helps prevent blood clotting and modulates bone health [24].

#### 2.2. Carotenoids

Carotenoids are tetraterpenoid lipids that can be divided into two major classes according to their chemical structure: carotenes that are nonoxygenated (e.g.,  $\alpha$ -carotene,  $\beta$ -carotene, and lycopene) and xanthophylls (e.g., lutein, zeaxanthin, fucoxanthin, and  $\beta$ -cryptoxanthin) that are oxygenated (Figure 1a). Carotenoids are abundant in yellow/orange fruit and vegetables, dark leafy green vegetables, and algae.  $\beta$ -carotene has pro-vitamin A function, since it is converted into vitamin A in the human body through metabolic pathways. Lutein and zeaxanthin retard the progression of age-related cataracts and help to improve cognitive function and cardiovascular health [2,25]. Lycopene has been reported to inhibit breast and prostate cancer, reduced the risk of cardiovascular disease, and retard nerval degenerative diseases [26]. Fucoxanthin can be isolated from seaweed and is claimed to exhibit anticancer, antidiabetic, antioxidant, anti-inflammatory, and antimicrobial activities, was as providing protection against liver, brain, bone, skin, and eye diseases [27,28].

## 2.3. Phytosterols

Phytosterols (plant sterols and stanols) are important structural components of plant cell membranes (Figure 1a) [12]. Sterols can occur in free form or esterified and glycosylated bound form. The most common forms of phytosterols found in foods are  $\beta$ -sitosterol, stigmasterol and campesterol. Phytosterols have been reported to reduce the risk of inflammatory diseases, as well as arteriosclerosis and cardiovascular diseases by lowering serum LDL-cholesterol concentrations [3,29].

#### 2.4. Essential Fatty Acids (Omega-3s and CLA)

Alpha-linolenic acid (ALA), eicosapentaenoic acid (EPA), and docosahexaenoic acid (DHA) are the major kinds of omega-3 polyunsaturated fatty acids (PUFAs) found in foods (Figure 1b) [11]. Fish oil is the main source of EPA and DHA in the human diet, although algae oil is also an important source of DHA [30]. Consumption of adequate amounts of omega-3 PUFAs has been reported to reduce the risk of cardiovascular diseases, boost neurological and visual functions, promote healthy aging, and improve cognitive functions in Alzheimer patients [4]. ALA is derived from plant seed oils and can be converted to EPA and DHA in vivo, but at a relatively low conversion rate [31]. Consequently, there is interest in fortifying plant-based foods with ALA as a non-animal source of healthy lipids [32]. Conjugated linoleic acids (CLA) are a group of positional and geometric isomers of linoleic acid. The most common bioactive isomers are cis-9, trans-11-CLA (9-CLA) and trans-10, cis-12-CLA (10-CLA), which belong to the n-7 and n-6 family, respectively. In vitro and animal models have shown that CLAs have various potential health benefits, including hypolipidemic, anti-carcinogenic, ostheosynthetic, anti-diabetagenic, and immunomodulatory effects. The major dietary sources of CLA are dairy and meat products from ruminants [33].

## 3. Nanoscale Delivery Systems for Encapsulation of Bioactive Lipids

Nanoscale delivery systems can be used to improve the handling, stability, and bioavailability of bioactive lipids [16,19,34,35]. For specific applications, nanotechnology has advantages over conventional encapsulation technologies, due to the smaller particle size and greater surface area of the carrier particles employed. Several lipid-based nanoscale delivery systems have been developed and to protect and deliver bioactive lipids, which differ in their compositions, structures, and functionalities (Figure 2) [16,36]. The functional performance of many of these delivery systems can be tailored for specific applications by changing the size, shape, charge, composition, or aggregation state of the nanoparticles they contain (Figure 3). A number of the most widely used nanoscale delivery systems for bioactive lipids are highlighted here.



**Figure 2.** Various kinds of edible nanoparticles are available to encapsulate, protect, and delivery bioactive substances in foods.



**Figure 3.** The properties of edible nanoscale delivery systems can be controlled to tailor their functionality.

## 3.1. Nanoemulsions

Oil-in-water nanoemulsions consist of small emulsifier-coated oil droplets (d < 200 nm) dispersed in water [37] (Figure 2). The bioactive lipids are typically trapped inside the oil droplets, which can be achieved either before or after nanoemulsion formation. This kind of nanoemulsion may be fabricated from food-grade oil, water, and emulsifier using low-or high-energy approaches. Low-energy approaches rely on the spontaneous formation of small oil droplets when the system composition or environmental conditions are changed in a specific fashion. High-energy approaches utilize mechanical devices ("homogenizers") to break up the oil and water phases and form small oil droplets that are coated by emulsifiers. Numerous food grade emulsifiers can be used to create edible nanoemulsions, including synthetic surfactants, biosurfactants, phospholipids, proteins, and polysaccharides [38–41]. Typically, the emulsifier-coated oil droplets are prevented from aggregating (flocculating or

coalescing) with each other by generating strong electrostatic and/or steric repulsive forces between them. The small size of the droplets in nanoemulsions provides strong resistance to gravitational separation [37]. However, nanoemulsions are susceptible to droplet growth through Ostwald ripening when the oil phase used has an appreciable solubility in water (such as for essential or flavor oils). In this case, they have to be carefully designed to reduce this form of instability, e.g., by adding a ripening inhibitor.

Nanoemulsions should be clearly distinguished from microemulsions, which are also used as bioactive delivery systems and have some similar compositional and structural characteristics [37]. However, nanoemulsions are thermodynamically unstable systems, whereas microemulsions are thermodynamically stable ones. As a result, microemulsions can often be formed by simply mixing bioactive, surfactant, and water together, whereas nanoemulsions require specific preparation methods (often involving high energy input). One drawback of microemulsions is that they can usually only be formulated from high concentrations of synthetic surfactants, which is undesirable for many food applications [42].

## 3.2. Solid Lipid Nanoparticles and Nanostructured Lipid Carriers

Solid lipid nanoparticles (SLN) also contain small emulsifier-coated lipid particles (d < 200 nm) but in this case the molecules within the lipid phase are organized into highly regular crystalline structures, rather than being disorganized as in a fluid [43] (Figure 2). The bioactive lipids are again located inside the lipid phase. If they are successfully trapped within the crystal structure formed by the lipid molecules, then they may have improved resistance to chemical degradation and a sustained release rate. However, they are sometimes expelled to the surfaces of the SLNs when the lipid phase crystallizes, which can actually reduce their chemical stability. Moreover, the lipid nanoparticles may change shape from spherical to non-spherical after formation, which can increase their surface area susceptibility to aggregation. For these reasons, the nature of both the lipid phase and emulsifier used must be carefully selected to ensure good performance of SLNs. Nanostructured lipid carriers (NLCs) were designed to overcome the problems with SLNs (Figure 2). They again consist of small emulsifier-coated lipid particles (d < 200 nm) dispersed in water, but in this case a mixture of solid and liquid lipids is used to create a less ordered lipid phase, which allows higher solubilization and retention of the bioactive lipids, as well as decreasing changes in particle shape during lipid phase crystallization [44,45].

#### 3.3. Nanoliposomes

Nanoliposomes are colloidal dispersions containing small (d < 200 nm) lipid nanoparticles comprised of one or more concentric lipid bilayers formed by self-association of phospholipid molecules in water [46] (Figure 2). At the particle surfaces, the polar head groups of the phospholipids protrude into the surrounding water phase. Within the concentric bilayers, the non-polar phospholipid tails face each other, which allows bioactive lipids to be trapped within these hydrophobic domains [34,35,47]. Nanoliposomes may be formed by various methods, including coating-solvent evaporation and homogenization methods [34], with the former being mainly used in the laboratory and the latter being more suitable for large-scale continuous commercial production.

#### 3.4. Biopolymer Nanogels and Nanofibers

Biopolymer nanogels and nanofibers have been developed to encapsulate bioactive lipids [35] (Figure 2). Nanogels consist of small biopolymer-rich particles (<200 nm) suspended in water. The particles contain a 3D network of crosslinked biopolymer molecules that trap a considerable amount of water through hydration and capillary forces. The biopolymer molecules may be crosslinked by physical or chemical interactions and the pore size of the network formed can be controlled by altering biopolymer or crosslinking concentration, as well as preparation conditions. Bioactive lipids may be directly encapsulated within the biopolymer network, or they may first be loaded into lipophilic nanocarriers

(such as nanoemulsions or nanoliposomes). Biopolymer nanogels can be designed to improve the stability and delivery of bioactive lipids [19,35]. Biopolymer nanofibers, which consist of thin fibers of crosslinked biopolymer molecules, are typically formed using electrospinning methods. Bioactive lipids can be directly or indirectly trapped within the biopolymer solution used to form the nanofibers prior to the electrospinning process. Nanochitin, nanocellulose, and nano-starch delivery systems have been produced using this method for food processing and packaging applications [48,49].

## 3.5. Nanoparticle-Stabilized Pickering Emulsions

Oil-in-water Pickering emulsions consist of oil droplets coated by colloidal particles [50,51] (Figure 2). These types of structured emulsions can be produced entirely from food grade ingredients and can be used to encapsulate bioactive lipids. Food grade nanoparticles that are suitable for used as emulsifiers in Pickering emulsions include nanocellulose, nanochitin, nano-starch, protein nanoparticles, and SLNs [51–54]. The bioactive lipids can be encapsulated within the oil droplets or within the nanoparticles used to stabilize them.

## 3.6. Nanoparticle Formation Methods

Nanoscale delivery systems can be produced using various low- or high-energy approaches, depending on the nature of the bioactive to be encapsulated, the nature of the encapsulating materials used to assemble the particles, and the intended functional attributes of the final system. High-energy methods include microfluidization, sonication, homogenization, and high-speed shearing methods, while low-energy methods include spontaneous emulsification, phase inversion, antisolvent precipitation, and evaporation methods [16,34,44]. Nanoemulsions, SLNs, and NLCs are commonly produced using both high-energy (sonication, homogenization, microfluidizer) and low-energy (spontaneous emulsification, phase inversion) methods. In the case of SLNs and NLCs, a nanoemulsion is typically prepared at high temperatures, which is then cooled below the melting point of the lipid phase [43]. Nanoliposomes can also be produced using both low- and highenergy methods, such as coating-evaporation and microfluidization approaches [16]. Other technologies, such as nano-spray drying and freeze drying can be used to create powders containing nanoparticles loaded with bioactive lipids [55–59]. Each nanoscale delivery system has its own pros and cons, which should be considered when selecting one for a specific food application.

#### 4. Improved Food Matrix Compatibility and Handling

Encapsulation of bioactive lipids in nanoparticles is widely used to improve their compatibility with different kinds of food matrices. The exterior of these nanoparticles is usually hydrophilic, which means they can be readily dispersed into the aqueous environments found in many foods and beverages, such as soft drinks, milk analogs, yogurts, dressings, sauces, and condiments. Nevertheless, the nanoparticles must also be designed to remain stable under the specific solution conditions found within a particular food product, such as its pH, ionic strength, and ingredient formulation. This means that any repulsive interactions between the nanoparticles, typically electrostatic and/or steric, must be designed to be stronger than any attractive interactions, typically van der Waals, hydrophobic, bridging and/or depletion [60]. This can be achieved by controlling the interfacial composition and structure, particularly the charge, thickness, and polarity of the nanoparticle coating.

Nanotechnology can also be used to improve the handling and storage of bioactive lipids. For instance, the bioactive lipids can be converted into a liquid, gel, powdered, or solid form that can facilitate their transport and utilization during the manufacturing process. Moreover, trapping the bioactive lipids within nanoparticles can improve their resistance to creaming, sedimentation, flocculation, or coalescence during storage, as well as their resistance to chemical degradation, thereby increasing the shelf-life of the food product [37]. Nevertheless, nanoscale delivery systems must be carefully designed to

retain the encapsulated bioactive lipids throughout storage, as well as to protect them from the specific environmental stressors they are exposed to, such as pH changes, heating, cooling, shearing, and dehydration. In the remainder of this section, examples of some of the nanoparticle-based systems used to improve the food matrix compatibility or handling of bioactive lipids are given.

Nanoliposomes have been used to incorporate fish oil into yogurts so as to fortify them with health-promoting omega-3 fatty acids [61]. Nanoencapsulation of the fish oils significantly increased the overall acceptance and taste, aroma, texture, and appearance of the yogurts compared utilization of non-encapsulated (free) fish oil. Fish oil has also been encapsulated within caseinate-gum arabic nanocomplexes that were converted into a powdered form and then introduced into fruit juices [62]. Sensory analysis of these omega-3 enriched fruit juices (60 mg EPA+DHA/100 mL) indicated an acceptable taste by consumer panelists, suggesting that nanoencapsulation masked off-flavors from the fish oil. In another study, fish oil was encapsulated within nanoliposomes that were then incorporated into bread, without adversely affecting the texture or sensory acceptability [63]. Conjugated linoleic acid (CLA) has been encapsulated within NLCs that were then added to low-fat milk [64]. Encapsulation was shown to improve the oxidative stability of the CLA, thereby extending the shelf life of the product.

Nanoencapsulation has also been used to improve the food matrix compatibility of various kinds of carotenoids. For instance, astaxanthin has been encapsulated within oil-in-water nanoemulsions that were then successfully introduced into orange juice and milk to form stable carotenoid-fortified products [65]. Zeaxanthin has been encapsulated in biopolymer nanoparticles and nanoemulsions and then incorporated into yogurt [66]. Encapsulation of the carotenoids improved their chemical stability, while having no adverse effects on the appearance, color, flavor, consistency, and texture of the fortified yogurts. Carotenoids extracted from Cantaloupe melon have also been encapsulated in nanoemulsions stabilized by whey proteins and gelatin, which were then incorporated into yogurt [67]. In this case, nanoencapsulation improve the dispersibility and color stability of the carotenoids in the yogurt compared to a crude carotenoid extract. Lycopene has been encapsulated within SLN and NLC systems, which were then successfully introduced into an orange drink without adversely affecting its sensory attributes [68].

Powdered forms of bioactive-loaded nanoparticles can be produced by spray drying, which improves the handling and storage stability. The powders produced can be directly incorporated into dried food products, such as soup mixes, fruit powders, cereal powders, and infant formulas, by simple mixing, or they can be redispersed in water and then incorporated into wet food products, such as beverages, soups, or sauces [69]. Flaxseed oil (56%  $\alpha$ -linolenic acid) has been encapsulated within nanocomplexes formulated from high amylose corn starch, which were then converted into a powdered form using spray drying [57]. The resulting powder was then used to fortify bread with omega-3 fatty acids. The encapsulated flaxseed oil had higher oxidative stability than free flaxseed oil during baking, which reduced off-flavor and acrylamide formation.

Vitamin  $D_3$  has been encapsulated within NLCs, which were then introduced into a yogurt-based beverage ("Lassi"), without adversely affecting the sensory attributes of the fortified product, such as appearance, flavor, aftertaste, and homogeneity [70]. Plantbased nanoemulsions have been used to encapsulate vitamin D and introduce them into plant-based milk analogs, such as almond and oat milks [71]. The nanoemulsions did not negatively alter the color or viscosity of the fortified milk analogs. Some examples of food and beverages that have been fortified with nanoencapsulated bioactive lipids are provided in Table 1.

<b>Bioactive Lipid</b>	Nanocarrier System	Food Application	Benefits	Reference
Polyunsaturated omega-3 and omega-6 fatty acids				
Flax seed oil (high in ALA)	Starch nanocomplexes	Bread	Reduced lipid oxidation, reduced HMF and acrylamide during baking	[57]
Fish oil (EPA-DHA)	Nanoliposomes	Yogurt	Increased bioactive retention, reduced lipid oxidation, high sensory score	[61]
Fish oil (EPA-DHA)	Sodium caseinate/gum arabic nanocomplexes	Fruit juice	Acceptable taste scores, high bioaccessibility	[62]
Fish oil (EPA-DHA)	Nanoliposomes	Bread	Improved oxidative stability, good textural and sensory quality	[63]
CLA	Nanostructured lipid carriers	Low fat milk	Improved physical, oxidative and thermal stability	[64]
		Carotenoids		
Astaxanthin	Nanodispersions	Orange juice, skimmed milk	Improved storage stability, high in vitro cellular uptake	[65]
Zeaxanthin	Nanoparticles, nanoemulsions	Yogurt	High storage stability in food matrix, in vitro controlled release	[66]
Carotenoid extract (high in β-carotene)	Nanoemulsion	Yogurt	Increased dispersibility, high storage stability	[67]
Lycopene	Solid lipid nanoparticles, nanostructured lipid carriers	Orange drink	Increased dispersibility, better aftertaste scores, high overall acceptance	[68]
Lycopene	Nanoemulsion	Model beverage	Improved storage stability	[72]
Vitamin D <sub>3</sub>	Nanostructured lipid carrier	Fat soluble vitamins 'Lassi' Yogurt based beverage	High sensory acceptance, sustained release in simulated intestinal fluid	[70]
Vitamin D <sub>3</sub>	Organic nanoparticles (Nanoemulsion, Nanocellulose),	Plant-based milks	Improved viscosity and color	[71]
Vitamin A, D <sub>3</sub> Vitamin D <sub>2</sub> Vitamin E	inorganic nanoparticle (TiO <sub>2</sub> ) Reassembled casein micelle Na-caseinate nanocomplexes Nanoemulsion	Skim milk Milk Fruit juice	Improved storage stability Improved storage and thermal stability Increased shelf life	[73] [74] [75]
	Phytosterols			
β-sitosterol	Nanostructured lipid carriers	Butter	Improved storage stability, high antioxidant activity	[76]
Phytosterols	Nanoporous starch aerogels	Granola bars, pudding	Improved bioaccessibility	[77]

Table 1. Effect of nanoencapsulation on applications of bioactive lipids in foods and beverages.

 $ALA: \alpha-linolenic \ acid, \ EPA: eicosapentaenoic \ acid, \ DHA: \ docosahexaenoic \ acid, \ CLA: \ conjugated \ linoleic \ acid.$ 

#### 5. Enhanced Stability

It is important that bioactive lipids remain stable after being incorporated into delivery systems or into fortified foods. A number of bioactive lipids are chemically unstable and so may degrade during the processing or storage of fortified foods, i.e., omega-3 fatty acids, CLA, vitamin E, and carotenoids [17]. These instability reactions are often accelerated by exposure to heat, oxygen, light, or pro-oxidants. In some cases, the chemical instability of bioactive lipids reduces their biological efficacy. In other cases, labile bioactive lipids can be used as natural antioxidants to prevent the oxidation of other labile bioactive lipids, e.g., carotenoids can be used as antioxidants for omega-3 fatty acids. As well as their chemical instability, it is also important to inhibit the physical instability of bioactive lipids within foods during processing, storage, and utilization, e.g., due to aggregation or gravitational separation.

Nanoencapsulation can be used to overcome problems associated with the chemical and physical stability of bioactive lipids. Vitamin A has been encapsulated within SLNs and NLCs using hot homogenization, which was shown to produce a physically and chemically stable form of these bioactive lipids [78]. Vitamin D<sub>3</sub> has been encapsulated within protein-polysaccharide nanocomplexes [79], which were shown to remain physically and chemically stable for 31 days when stored at 4 °C. In another study, it was shown that encapsulation of vitamin A in caseinate nanocomplexes improved its storage stability compared to free vitamin A [80]. The stability of vitamin D<sub>2</sub> in milk to pasteurization, boiling, and sterilization improved after it was encapsulated within caseinate nanocomplexes [74].

Moreover, the stability of the encapsulated vitamin  $D_2$  was higher than that of free vitamin  $D_2$  during 6-months storage of the fortified milks at -20, 4, and 37 °C, as well as when the milks were exposed to light. Nanoemulsions fortified with vitamin  $D_3$  were shown to remain relatively stable when exposed to a range of environmental stresses, including heating, chilling, pH variations (pH 3 to 7), and ionic strength (0–750 mM) [81]. The highest retention was achieved under refrigerated conditions, at lower ionic strengths, and at pH 7.0. Encapsulation within nanoemulsions has also been shown to improve the oxidative stability of vitamin E during long-term storage [82]. Vitamin E encapsulated within nanoemulsions has been used to fortify mango juice [75], leading to a product that had good physical and chemical stability during storage. Nanoemulsions have also been shown to improve the stability of vitamin E to thermal degradation when exposed to various heat treatments, i.e., 30–90 °C for 30 min [83]. Rapid degradation of vitamin E was only observed at the highest temperature used (90 °C). The vitamin E nanoemulsions exhibited good long-term stability (60 days) when stored in the light and in the dark at 4 and 25  $^{\circ}$ C, but some degradation was observed during storage at 40 °C. The authors also showed that the storage stability of the vitamin E was higher when long-chain triglycerides were used as an oil phase rather than shorter chain ones. Nanoemulsions have also been shown to improve the chemical stability of vitamin E during 4-weeks storage in the dark [84]. Reducing the oil droplet size in these nanoemulsions also improved the storage stability of the nanoemulsions to gravitational separation.

The stability of  $\beta$ -carotene encapsulated within nanoemulsions has been examined when they were exposed to thermal treatments (pasteurization and sterilization), as well as they were stored in the light and the dark at 4, 25, and 37 °C for up to 90 days [85]. Storage at 4 °C in the dark was found to be the best conditions to prevent  $\beta$ -carotene degradation.

Encapsulation of lutein within nanoemulsions stabilized by food grade emulsifiers has been shown to improve the chemical stability of the carotenoid in nutraceutical-fortified beverages [86]. The physical stability of the nanoemulsions to aggregation and creaming could be controlled by selecting an appropriate emulsifier. The physical and chemical stability of lycopene encapsulated in nanoemulsions has been determined when they were stored at 4, 20, 30 and 40 °C for 3 weeks in the dark [72]. The results showed that the rate of lycopene degradation decreased as the temperature decreased.

DHA has been successfully encapsulated within casein micelles (r-CM) and casein nanoparticles (CNP), which improved the chemical stability of the polyunsaturated lipids during storage at 4 °C for 16 and 37 days [87]. DHA has also been encapsulated within  $\beta$ -lactoglobulin-pectin nanodispersions, which improved its resistance to oxidation when exposed to an accelerated shelf-life stress test, i.e., 100 h at 40 °C [88]. Indeed, there was only 5–10% degradation of DHA in the encapsulated form compared to 80% for the bulk form. The encapsulation of krill oil, which is high in omega-3 fatty acids, within nanoemulsions upon the stability of the oil to oxidation during freeze drying, spray drying, and storage for 12 days has been examined [58]. Encapsulation in the nanoemulsions actually reduced the oxidative stability of the krill oil, which was attributed to the increased exposure of the oil phase to air as a result of the increase in surface area. The addition of a natural antioxidant (vitamin E) has been shown to improve the oxidative stability of ALA encapsulated within nanoemulsions [89]. In this case, the encapsulated ALA had a higher oxidative stability than the free form. Higher temperatures and the presence of metal ions increased ALA degradation, whereas light exposure and ionic strength did not significantly change ALA degradation.

The physical and chemical stability of omega-3,  $\beta$ -carotene and vitamin A encapsulated within SLNs has been studied during 51-days storage [17]. The physical stability of the bioactive-loaded SLNs could be improved by using suitable surfactants, such as *quillaja saponin* and its combination with high-melting lecithin. The oxidative stability of the encapsulated bioactive lipids was improved by preventing the polymorphic transition of the lipid phase in the SLNs by using high-melting surfactants. The oxidative stability of EPA and DHA was improved after encapsulation within nanoliposomes [90], while that of ALA and quercetin was improved after co-encapsulation in NLCs [91]. Encapsulation of bixin (a carotenoid) within nanocapsules was found to improve its chemical stability when exposed to light, oxygen, and heat [92]. A phytosterol ( $\beta$ -sitosterol) has been encapsulated within NLCs, which were then incorporated into butter [76]. Fortification of the butter with  $\beta$ -sitosterol increased its resistance to oxidation during long-term storage (3-months) in a refrigerator.

Pickering emulsions stabilized with food-grade nanoparticles have been reported to be good candidates for use in processed foods due to their higher stability against environmental stresses such as freeze-thawing, extreme pH conditions, and high ionic strengths [50]. Pickering emulsions stabilized by chitosan hydrochloride/carboxymethyl starch nanogels have been used to encapsulate  $\beta$ -carotene, which improved the stability of the carotenoid to UV exposure and heating [93]. Pickering emulsions stabilized by egg yolk peptides and micellar nanoparticles were found to be stable to spray drying conditions and to oil leakage during storage [54]. Similarly, encapsulation of  $\beta$ -carotene in Pickering emulsions stabilized by gliadin nanoparticles and gum arabic has been shown to improve the resistance of the carotenoid to pH, salt, heat, and light [94]. Pickering emulsions have also been used to improve the chemical stability of astaxanthin [95]. In this case, the Pickering emulsions were stabilized by lupin protein aggregates.

## 6. Enhanced Bioavailability

The bioavailability of bioactive substances is influenced by their liberation, solubilization, absorption, and metabolism within the gastrointestinal tract (GIT) (Figure 4) [96]. For lipophilic bioactive substances, the rate limiting steps determining their bioavailability is often their poor solubilization within intestinal fluids, as well as their chemical degradation. The bioavailability of these lipophilic bioactives can therefore be improved by increasing their bioaccessibility and inhibiting their degradation within the gastrointestinal tract [96]. Nanoencapsulation has been widely used to improve the bioavailability of this type of bioactive substance [18].



**Figure 4.** The bioavailability of hydrophobic bioactive substances can be increased by encapsulating them in nanoparticles. The triglycerides (TG) in the lipid nanoparticles are broken down to free fatty acids (FFA) and monoglycerides (MG), which are packed into mixed micelles with bioactive substances and transported to the epithelium cells. They are then reassembled into triglycerides, packed into chylomicrons, and transported into the bloodstream through the lymphatic system. Lipid digestion and nutraceutical bioaccessibility increases with decreasing droplet size.

Lipid nanoparticles undergo a number of disassembly and assembly processes inside the human body that impacts the bioaccessibility of any hydrophobic bioactives encapsulated within them [97]. The nanoparticles are digested by lipases in the stomach and small intestine, which releases free fatty acids and monoacylglycerols that are incorporated into mixed micelles with bile salts and phospholipids. These mixed micelles then solubilize the hydrophobic bioactives and transport them to the epithelial cells where they can be absorbed. After absorption, the hydrophobic bioactives are packed into chylomicrons that carry them into the bloodstream via the lymphatic system, thereby by-passing first-pass metabolism within the liver [98]. The majority of bioavailability studies have been carried out using in vitro simulated gastrointestinal tract (GIT) models due to their relatively low cost, simplicity, and speed. However, researchers may also use Caco-2 cell lines or animal models to study the bioavailability of bioactive lipids. In particular, co-cultured cell lines containing different kinds of cells (such as model enterocytes, M-cells, and mucus-secreting cells) are particularly useful for mimicking the lining of the human GIT [99–101].

As an example, omega-3 fatty acids loaded into nanoemulsions have been shown to increase the bioaccessibility of EPA and DHA compared to bulk oil using a simulated GIT model [102]. Encapsulation of astaxanthin in potato protein-based nanoparticles was shown to improve the bioaccessibility, chemical stability, and bioaccessibility of the carotenoids under simulated gastrointestinal conditions [103]. For instance, encapsulation increased the bioaccessibility by 11-fold and the bioavailability by 7-fold.

The encapsulation of vitamin A into caseinate nano-complexes has been shown to increase the in vitro bioaccessibility and bioavailability of this fat soluble vitamin [104]. In particular, nanoencapsulation protected the vitamin A from degradation under the acidic conditions in the simulated stomach, as well as increasing the fraction absorbed by Caco-2 cells compared to the free form. NLCs have been used to increase the bioavailability of vitamin D<sub>3</sub>, which again was partly attributed to its ability to increase the chemical stability under simulated gastric conditions [105]. Similarly, encapsulation of vitamin D<sub>3</sub> in ovalbumin-pectin nanocomplexes has been shown to protect the vitamin from degradation under gastric conditions [79].

Oil-in-water nanoemulsions stabilized by a natural surfactant (quillaja saponin) have been shown to increase the bioaccessibility of vitamin  $D_3$  [106]. The nature of the oil phase has been shown to have a major impact on the bioaccessibility of lipophilic bioactives. For instance, vitamin E encapsulated within nanoemulsions formulated from long-chain triglycerides (LCT) had a higher bioaccessibility than those formulated from medium-chain triglycerides (MCT) [107]. In another study, encapsulation of gamma- and delta-tocotrienols in nanoemulsions was shown to increase their bioaccessibility compared to conventional emulsions using a simulated GIT [108]. In addition, encapsulation of vitamin E in plantbased nanoemulsions was shown to increase its bioaccessibility by an amount that depends on oil phase composition [84].

Encapsulation of lutein within sophorolipid-coated zein nanoparticles has been shown to improve the bioaccessibility of the carotenoid using a GIT model [109]. Similarly, encapsulation of lycopene into oil-in-water nanoemulsions increased carotenoid bioaccessibility, with the efficacy increasing with decreasing droplet size [110]. Phytosterols have been encapsulated within pectin-coated zein nanoparticles, which was shown to improve their stability and to modulate their release under simulated GIT conditions [111]. Phytosterols have also been encapsulated within nanoporous starch aerogels [77], which increased their in vitro bioaccessibility when incorporated into granola bars and puddings compared to free phytosterols.

Excipient foods are designed to increase the bioavailability of bioactive substances in other foods that are co-ingested with them. Excipient nanoemulsions have been used to enhance the oral bioavailability of various kinds of bioactive lipids, which has been attributed to their ability to increase the bioaccessibility, retard the chemical degradation, and increase the absorption of these compounds [112]. For instance, oil-in-water nanoemulsions have been used to increase the bioaccessibility of lycopene in tomato juice [113]. This effect was attributed to the rapid formation of mixed micelles in the small intestine, which could solubilize the lycopene. Nanoemulsions have also been used to increase the in vitro bioaccessibility and absorption of  $\beta$ -carotene using simulated GIT models, which was again attributed to increased mixed micelle formation [93,114]. Encapsulation of EPA and DHA within nanoemulsions has also been shown to increase their in vivo bioavailability using rat models [115]. In another study, the oral bioavailability of vitamin E ( $\gamma$ -tocotrienol) was enhanced three-fold when orally administered to rats in the form of SLNs [116]. Overall, all of these studies show that various kinds of edible nanoemulsions can be used to increase the bioavailability of bioactive lipids.

#### 7. Conclusions

There are a wide variety of different bioactive lipids found in foods that may exhibit health benefits in humans. However, these benefits may not be fully realized because many bioactive lipids are difficult to incorporate into foods, are degraded in foods during storage, or have low bioavailability after ingestion. Nanotechnology can be used to improve the matrix compatibility, physiochemical stability, and bioavailability of bioactive lipids. In particular, edible nanoparticles can be created from food grade ingredients to overcome these challenges in food applications. This review has highlighted the characteristics of different kinds of bioactive lipids (e.g., oil-soluble vitamins, carotenoids, phytosterols, and essential fatty acids), as well as the different types of nanoscale delivery systems that can be used to encapsulate them (such as nanoemulsions, SLNs, nanoliposomes, biopolymer nanogels, and Pickering emulsions). In the future, it will be important to identify the most appropriate nanoscale delivery systems for each application, to test the optimized delivery systems using in vivo animal and human studies, and to ensure that the delivery systems remain viable within real food applications.

**Author Contributions:** Conceptualization, D.J.M. and B.Ö.; writing—original draft preparation, B.Ö.; writing—review and editing, D.J.M.; visualization, D.J.M. and B.Ö.; supervision, D.J.M. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by the National Institute of Food and Agriculture, USDA, Massachusetts Agricultural Experiment Station (Project Number 831) and USDA, AFRI (2020-03921) Grants.

Acknowledgments: This material was partly based upon work supported by the National Institute of Food and Agriculture, USDA, Massachusetts Agricultural Experiment Station (Project Number 831) and USDA, AFRI (2020-03921) Grants.

Conflicts of Interest: The authors declare no conflict of interest.

#### References

- Sankar, R.; Ravisankar, P.; Reddy, A.A.; Nagalakshmi, B.; Koushik, O.S.; Vijaya Kumar, B.; Anvith, P.S. The comprehensive review on fat soluble vitamins. *IOSR J. Pharm.* 2015, 5, 12–28.
- 2. Eggersdorfer, M.; Wyss, A. Carotenoids in Human Nutrition and Health. Arch. Biochem. Biophys. 2018, 652, 18–26. [CrossRef]
- 3. Trautwein, E.A.; Demonty, I. Phytosterols: Natural Compounds with Established and Emerging Health Benefits. *Oléagineux Corps Gras Lipides* 2007, 14, 259–266. [CrossRef]
- 4. Swanson, D.; Block, R.; Mousa, S.A. Omega-3 Fatty Acids EPA and DHA: Health Benefits throughout Life. *Adv. Nutr.* **2012**, *3*, 1–7. [CrossRef]
- Chen, B.; McClements, D.J.; Decker, E.A. Design of Foods with Bioactive Lipids for Improved Health. Annu. Rev. Food Sci. Technol. 2013, 4, 35–56. [CrossRef]
- Vázquez, L.; Corzo-Martínez, M.; Arranz-Martínez, P.; Barroso, E.; Reglero, G.; Torres, C. Bioactive Lipids. In *Bioactive Molecules in Food. Reference Series in Phytochemistry*; Mérillon, J.M., Ramawat, K.G., Eds.; Springer International Publishing: Cham, Switzerland, 2019; pp. 467–527.
- 7. Das, U.N. Bioactive Lipids in COVID-19-Further Evidence. Arch. Med. Res. 2020. [CrossRef]
- Rogero, M.M.; Leão, M.D.C.; Santana, T.M.; Pimentel, M.V.D.M.; Carlini, G.C.; Da Silveira, T.F.; Gonçalves, R.C.; Castro, I.A. Potential Benefits and Risks of Omega-3 Fatty Acids Supplementation to Patients with COVID-19. *Free Radic. Biol. Med.* 2020, 156, 190–199. [CrossRef] [PubMed]
- 9. Bistrian, B.R. Parenteral Fish-Oil Emulsions in Critically Ill COVID-19 Emulsions. J. Parenter. Enter. Nutr. 2020, 44, 1168. [CrossRef] [PubMed]
- Meltzer, D.O.; Best, T.J.; Zhang, H.; Vokes, T.; Arora, V.; Solway, J. Association of Vitamin D Status and Other Clinical Characteristics with COVID-19 Test Results. JAMA Netw. Open 2020, 3, e2019722. [CrossRef]

- Saini, R.K.; Keum, Y.S. Omega-3 and Omega-6 Polyunsaturated Fatty Acids: Dietary Sources, Metabolism, and Significance—A Review. *Life Sci.* 2018, 203, 255–267. [CrossRef]
- Moreau, R.A.; Nyström, L.; Whitaker, B.D.; Winkler-Moser, J.K.; Baer, D.J.; Gebauer, S.K.; Hicks, K.B. Phytosterols and Their Derivatives: Structural Diversity, Distribution, Metabolism, Analysis, and Health-Promoting Uses. *Prog. Lipid Res.* 2018, 70, 35–61. [CrossRef]
- 13. Maiani, G.; Castón, M.J.P.; Catasta, G.; Toti, E.; Cambrodón, I.G.; Bysted, A.; Granado-Lorencio, F.; Olmedilla-Alonso, B.; Knuthsen, P.; Valoti, M.; et al. Carotenoids: Actual Knowledge on Food Sources, Intakes, Stability and Bioavailability and Their Protective Role in Humans. *Mol. Nutr. Food Res.* **2009**, *53*, 194–218. [CrossRef]
- Tabibian, M.; Torbati, M.; Mogaddam, M.R.A.; Mirlohi, M.; Sadeghi, M.; Mohtadinia, J. Evaluation of Vitamin D3 and D2 Stability in Fortified Flat Bread Samples during Dough Fermentation, Baking and Storage. *Adv. Pharm. Bull.* 2017, 7, 323–328. [CrossRef]
- 15. Dutta, D.; Chaudhuri, U.R.; Chakraborty, R. Structure, Health Benefits, Antioxidant Property and Processing and Storage of Carotenoids. *Afr. J. Biotechnol.* **2005**, *4*, 1510–1520. [CrossRef]
- 16. McClements, D.J. Nanoscale Nutrient Delivery Systems for Food Applications: Improving Bioactive Dispersibility, Stability, and Bioavailability. *J. Food Sci.* 2015, *80*, N1602–N1611. [CrossRef] [PubMed]
- Salminen, H.; Gömmel, C.; Leuenberger, B.H.; Weiss, J. Influence of Encapsulated Functional Lipids on Crystal Structure and Chemical Stability in Solid Lipid Nanoparticles: Towards Bioactive-Based Design of Delivery Systems. *Food Chem.* 2016, 190, 928–937. [CrossRef] [PubMed]
- Yao, M.; McClements, D.J.; Xiao, H. Improving Oral Bioavailability of Nutraceuticals by Engineered Nanoparticle-Based Delivery Systems. *Curr. Opin. Food Sci.* 2015, 2, 14–19. [CrossRef]
- Shin, G.H.; Kim, J.T.; Park, H.J. Recent Developments in Nanoformulations of Lipophilic Functional Foods. *Trends Food Sci. Technol.* 2015, 46, 144–157. [CrossRef]
- Albahrani, A.A.; Greaves, R.F. Fat-Soluble Vitamins: Clinical Indications and Current Challenges for Chromatographic Measurement. *Clin. Biochem. Rev.* 2016, 37, 27–47. [PubMed]
- 21. Clagett-Dame, M.; Knutson, D. Vitamin a in Reproduction and Development. Nutrients 2011, 3, 385–428. [CrossRef]
- Prietl, B.; Treiber, G.; Pieber, T.R.; Amrein, K. Vitamin D and Immune Function. *Nutrients* 2013, *5*, 2502–2521. [CrossRef] [PubMed]
  Borel, P.; Desmarchelier, C. Annual Review of Nutrition Bioavailability of Fat-Soluble Vitamins and Phytochemicals in Humans: Effects of Genetic Variation. *Annu. Rev. Nutr.* 2018. [CrossRef] [PubMed]
- 24. Vermeer, C. Vitamin K: The Effect on Health beyond Coagulation—An Overview. *Food Nutr. Res.* **2012**, *56*, 5329. [CrossRef] [PubMed]
- 25. Boon, C.S.; McClements, D.J.; Weiss, J.; Decker, E.A. Factors Influencing the Chemical Stability of Carotenoids in Foods. *Crit. Rev. Food Sci. Nutr.* **2010**, *50*, 515–532. [CrossRef] [PubMed]
- Liang, X.; Ma, C.; Yan, X.; Liu, X.; Liu, F. Advances in Research on Bioactivity, Metabolism, Stability and Delivery Systems of Lycopene. *Trends Food Sci. Technol.* 2019, 93, 185–196. [CrossRef]
- 27. Lourenço-Lopes, C.; Garcia-Oliveira, P.; Carpena, M.; Fraga-Corral, M.; Jimenez-Lopez, C.; Pereira, A.G.; Prieto, M.A.; Simal-Gandara, J. Scientific Approaches on Extraction, Purification and Stability for the Commercialization of Fucoxanthin Recovered from Brown Algae. *Foods* **2020**, *9*, 1113. [CrossRef]
- 28. Peng, J.; Yuan, J.P.; Wu, C.F.; Wang, J.H. Fucoxanthin, a Marine Carotenoid Present in Brown Seaweeds and Diatoms: Metabolism and Bioactivities Relevant to Human Health. *Mar. Drugs* **2011**, *9*, 1806–1828. [CrossRef]
- Hannan, M.A.; Sohag, A.A.M.; Dash, R.; Haque, M.N.; Mohibbullah, M.; Oktaviani, D.F.; Hossain, M.T.; Choi, H.J.; Moon, I.S. Phytosterols of Marine Algae: Insights into the Potential Health Benefits and Molecular Pharmacology. *Phytomedicine* 2020, 69, 153201. [CrossRef]
- 30. Ryan, L.; Symington, A.M. Algal-Oil Supplements Are a Viable Alternative to Fish-Oil Supplements in Terms of Docosahexaenoic Acid (22:6n-3; DHA). *J. Funct. Foods* **2015**, *19*, 852–858. [CrossRef]
- 31. Baker, E.J.; Miles, E.A.; Burdge, G.C.; Yaqoob, P.; Calder, P.C. Metabolism and Functional Effects of Plant-Derived Omega-3 Fatty Acids in Humans. *Prog. Lipid Res.* 2016, *64*, 30–56. [CrossRef]
- 32. Sugasini, D.; Lokesh, B.R. Uptake of α-Linolenic Acid and Its Conversion to Long Chain Omega-3 Fatty Acids in Rats Fed Microemulsions of Linseed Oil. *Lipids* **2012**, *47*, 1155–1167. [CrossRef]
- 33. Benjamin, S.; Spener, F. Conjugated Linoleic Acids as Functional Food: An Insight into Their Health Benefits. *Nutr. Metab.* 2009, *6*, 36. [CrossRef] [PubMed]
- 34. Akhavan, S.; Assadpour, E.; Katouzian, I.; Jafari, S.M. Lipid Nano Scale Cargos for the Protection and Delivery of Food Bioactive Ingredients and Nutraceuticals. *Trends Food Sci. Technol.* **2018**, 74, 132–146. [CrossRef]
- Assadpour, E.; Mahdi Jafari, S. A Systematic Review on Nanoencapsulation of Food Bioactive Ingredients and Nutraceuticals by Various Nanocarriers. Crit. Rev. Food Sci. Nutr. 2019, 59, 3129–3151. [CrossRef] [PubMed]
- Sivakamasundari, S.K.; Leena, M.; Moses, J.A.; Anandharamakrishnan, C. Solid Lipid Nanoparticles: Formulation and Applications in Food Bioactive Delivery. In *Innovative Food Processing Technologies*; Elsevier: Amsterdam, The Netherlands, 2021; pp. 580–604.
- 37. McClements, D.J.; Rao, J. Food-Grade Nanoemulsions: Formulation, Fabrication, Properties, Performance, Biological Fate, and Potential Toxicity. *Crit. Rev. Food Sci. Nutr.* 2011, *51*, 285–330. [CrossRef]

- Shu, G.; Khalid, N.; Chen, Z.; Neves, M.A.; Barrow, C.J.; Nakajima, M. Formulation and Characterization of Astaxanthin-Enriched Nanoemulsions Stabilized Using Ginseng Saponins as Natural Emulsifiers. *Food Chem.* 2018, 255, 67–74. [CrossRef] [PubMed]
- Ozturk, B.; McClements, D.J. Progress in Natural Emulsifiers for Utilization in Food Emulsions. Curr. Opin. Food Sci. 2016, 7. [CrossRef]
- 40. Komaiko, J.; McClements, D.J. Low-Energy Formation of Edible Nanoemulsions by Spontaneous Emulsification: Factors Influencing Particle Size. J. Food Eng. 2015, 146, 122–128. [CrossRef]
- 41. Qian, C.; McClements, D.J. Formation of Nanoemulsions Stabilized by Model Food-Grade Emulsifiers Using High-Pressure Homogenization: Factors Affecting Particle Size. *Food Hydrocoll.* **2011**, 25, 1000–1008. [CrossRef]
- 42. McClements, D.J. Nanoemulsions versus Microemulsions: Terminology, Differences, and Similarities. *Soft Matter* 2012, *8*, 1719–1729. [CrossRef]
- Weiss, J.; Decker, E.A.; McClements, D.J.; Kristbergsson, K.; Helgason, T.; Awad, T. Solid Lipid Nanoparticles as Delivery Systems for Bioactive Food Components. *Food Biophys.* 2008, *3*, 146–154. [CrossRef]
- 44. Katouzian, I.; Faridi Esfanjani, A.; Jafari, S.M.; Akhavan, S. Formulation and Application of a New Generation of Lipid Nano-Carriers for the Food Bioactive Ingredients. *Trends Food Sci. Technol.* **2017**, *68*, 14–25. [CrossRef]
- Tamjidi, F.; Shahedi, M.; Varshosaz, J.; Nasirpour, A. Nanostructured Lipid Carriers (NLC): A Potential Delivery System for Bioactive Food Molecules. *Innov. Food Sci. Emerg. Technol.* 2013, 19, 29–43. [CrossRef]
- 46. Khorasani, S.; Danaei, M.; Mozafari, M.R. Nanoliposome Technology for the Food and Nutraceutical Industries. *Trends Food Sci. Technol.* **2018**, *79*, 106–115. [CrossRef]
- 47. Tan, C.; Zhang, Y.; Abbas, S.; Feng, B.; Zhang, X.; Xia, S. Modulation of the Carotenoid Bioaccessibility through Liposomal Encapsulation. *Colloids Surf. B Biointerfaces* **2014**, *123*, 692–700. [CrossRef] [PubMed]
- 48. Senthil Muthu Kumar, T.; Senthil Kumar, K.; Rajini, N.; Siengchin, S.; Ayrilmis, N.; Varada Rajulu, A. A Comprehensive Review of Electrospun Nanofibers: Food and Packaging Perspective. *Compos. Part B Eng.* **2019**, *175*, 175. [CrossRef]
- 49. Rostamabadi, H.; Assadpour, E.; Tabarestani, H.S.; Falsafi, S.R.; Jafari, S.M. Electrospinning Approach for Nanoencapsulation of Bioactive Compounds; Recent Advances and Innovations. *Trends Food Sci. Technol.* **2020**, *100*, 190–209. [CrossRef]
- 50. Weiss, J.; Ahmad, T.; Zhang, C.; Zhang, H. A Review of Recent Progress on High Internal-Phase Pickering Emulsions in Food Science. *Trends Food Sci. Technol.* 2020, 106, 91–103. [CrossRef]
- 51. Mwangi, W.W.; Lim, H.P.; Low, L.E.; Tey, B.T.; Chan, E.S. Food-Grade Pickering Emulsions for Encapsulation and Delivery of Bioactives. *Trends Food Sci. Technol.* 2020, 100, 320–332. [CrossRef]
- 52. Ren, Z.; Chen, Z.; Zhang, Y.; Lin, X.; Li, B. Novel Food-Grade Pickering Emulsions Stabilized by Tea Water-Insoluble Protein Nanoparticles from Tea Residues. *Food Hydrocoll.* **2019**, *96*, 322–330. [CrossRef]
- 53. Zhou, H.; Tan, Y.; Lv, S.; Liu, J.; Muriel Mundo, J.L.; Bai, L.; Rojas, O.J.; McClements, D.J. Nanochitin-Stabilized Pickering Emulsions: Influence of Nanochitin on Lipid Digestibility and Vitamin Bioaccessibility. *Food Hydrocoll.* **2020**, *106*. [CrossRef]
- 54. Du, Z.; Li, Q.; Li, J.; Su, E.; Liu, X.; Wan, Z.; Yang, X. Self-Assembled Egg Yolk Peptide Micellar Nanoparticles as a Versatile Emulsifier for Food-Grade Oil-in-Water Pickering Nanoemulsions. *J. Agric. Food Chem.* **2019**, *67*, 11728–11740. [CrossRef]
- 55. Salminen, H.; Ankenbrand, J.; Zeeb, B.; Badolato Bönisch, G.; Schäfer, C.; Kohlus, R.; Weiss, J. Influence of Spray Drying on the Stability of Food-Grade Solid Lipid Nanoparticles. *Food Res. Int.* **2019**, *119*, 741–750. [CrossRef]
- Bejrapha, P.; Min, S.G.; Surassmo, S.; Choi, M.J. Physicothermal Properties of Freeze-Dried Fish Oil Nanocapsules Frozen under Different Conditions. Dry. Technol. 2010, 28, 481–489. [CrossRef]
- 57. Gökmen, V.; Mogol, B.A.; Lumaga, R.B.; Fogliano, V.; Kaplun, Z.; Shimoni, E. Development of Functional Bread Containing Nanoencapsulated Omega-3 Fatty Acids. J. Food Eng. 2011, 105, 585–591. [CrossRef]
- 58. El-Messery, T.M.; Altuntas, U.; Altin, G.; Özçelik, B. The Effect of Spray-Drying and Freeze-Drying on Encapsulation Efficiency, in Vitro Bioaccessibility and Oxidative Stability of Krill Oil Nanoemulsion System. *Food Hydrocoll.* **2020**, *106*. [CrossRef]
- Chen, C.C.; Wagner, G. Vitamin E Nanoparticle for Beverage Applications. *Chem. Eng. Res. Des.* 2004, *82*, 1432–1437. [CrossRef]
  McClements, D.J. *Food Emulsions*, 3rd ed.; CRC Press: Boca Raton, FL, USA, 2015; ISBN 9780429154034.
- 61. Ghorbanzade, T.; Jafari, S.M.; Akhavan, S.; Hadavi, R. Nano-Encapsulation of Fish Oil in Nano-Liposomes and Its Application in Fortification of Yogurt. *Food Chem.* **2017**, *216*, 146–152. [CrossRef] [PubMed]
- 62. Ilyasoglu, H.; El, S.N. Nanoencapsulation of EPA/DHA with Sodium Caseinate-Gum Arabic Complex and Its Usage in the Enrichment of Fruit Juice. *LWT Food Sci. Technol.* **2014**, *56*, 461–468. [CrossRef]
- 63. Ojagh, S.M.; Hasani, S. Characteristics and Oxidative Stability of Fish Oil Nano-Liposomes and Its Application in Functional Bread. *J. Food Meas. Charact.* 2018, 12, 1084–1092. [CrossRef]
- 64. Hashemi, F.S.; Farzadnia, F.; Aghajani, A.; Ahmadzadeh NobariAzar, F.; Pezeshki, A. Conjugated Linoleic Acid Loaded Nanostructured Lipid Carrier as a Potential Antioxidant Nanocarrier for Food Applications. *Food Sci. Nutr.* **2020**, *8*, 4185–4195. [CrossRef] [PubMed]
- 65. Anarjan, N.; Tan, C.P. Chemical Stability of Astaxanthin Nanodispersions in Orange Juice and Skimmed Milk as Model Food Systems. *Food Chem.* **2013**, *139*, 527–531. [CrossRef]
- 66. de Campo, C.; Queiroz Assis, R.; Marques da Silva, M.; Haas Costa, T.M.; Paese, K.; Stanisçuaski Guterres, S.; de Oliveira Rios, A.; Hickmann Flôres, S. Incorporation of Zeaxanthin Nanoparticles in Yogurt: Influence on Physicochemical Properties, Carotenoid Stability and Sensory Analysis. *Food Chem.* 2019, 301. [CrossRef]

- Medeiros, A.K.D.O.C.; de Carvalho Gomes, C.; de Araújo Amaral, M.L.Q.; de Medeiros, L.D.G.; Medeiros, I.; Porto, D.L.; Aragão, C.F.S.; Maciel, B.L.L.; de AraújoMorais, A.H.; SouzaPassos, T. Nanoencapsulation Improved Water Solubility and Color Stability of Carotenoids Extracted from Cantaloupe Melon (*Cucumis Melo* L.). *Food Chem.* 2019, 270, 562–572. [CrossRef]
- Akhoond Zardini, A.; Mohebbi, M.; Farhoosh, R.; Bolurian, S. Production and Characterization of Nanostructured Lipid Carriers and Solid Lipid Nanoparticles Containing Lycopene for Food Fortification. *J. Food Sci. Technol.* 2018, 55, 287–298. [CrossRef] [PubMed]
- 69. Chopde, S.; Datir, R.; Deshmukh, G.; Dhotre, A.; Patil, M. Nanoparticle Formation by Nanospray Drying & Its Application in Nanoencapsulation of Food Bioactive Ingredients. J. Agric. Food Res. 2020, 2, 100085. [CrossRef]
- 70. Maurya, V.K.; Aggarwal, M. Fabrication of Nano-Structured Lipid Carrier for Encapsulation of Vitamin D3 for Fortification of 'Lassi'; A Milk Based Beverage. J. Steroid Biochem. Mol. Biol. 2019, 193. [CrossRef]
- 71. Zhou, H.; Liu, J.; Dai, T.; Muriel Mundo, J.L.; Tan, Y.; Bai, L.; McClements, D.J. The Gastrointestinal Fate of Inorganic and Organic Nanoparticles in Vitamin D-Fortified Plant-Based Milks. *Food Hydrocoll.* **2021**, *112*. [CrossRef]
- 72. Kim, S.O.; Ha, T.V.A.; Choi, Y.J.; Ko, S. Optimization of Homogenization-Evaporation Process for Lycopene Nanoemulsion Production and Its Beverage Applications. *J. Food Sci.* 2014, 79. [CrossRef]
- 73. Loewen, A.; Chan, B.; Li-Chan, E.C.Y. Optimization of Vitamins A and D3 Loading in Re-Assembled Casein Micelles and Effect of Loading on Stability of Vitamin D3 during Storage. *Food Chem.* **2018**, 240, 472–481. [CrossRef]
- 74. Syama, M.A.; Arora, S.; Gupta, C.; Sharma, A.; Sharma, V. Enhancement of Vitamin D 2 Stability in Fortified Milk during Light Exposure and Commercial Heat Treatments by Complexation with Milk Proteins. *Food Biosci.* **2019**, *29*, 17–23. [CrossRef]
- 75. Dasgupta, N.; Ranjan, S.; Mundra, S.; Ramalingam, C.; Kumar, A. Fabrication of Food Grade Vitamin E Nanoemulsion by Low Energy Approach, Characterization and Its Application. *Int. J. Food Prop.* **2016**, *19*, 700–708. [CrossRef]
- 76. Bagherpour, S.; Alizadeh, A.; Ghanbarzadeh, S.; Mohammadi, M.; Hamishehkar, H. Preparation and Characterization of Betasitosterol-Loaded Nanostructured Lipid Carriers for Butter Enrichment. *Food Biosci.* **2017**, *20*, 51–55. [CrossRef]
- Ubeyitogullari, A.; Ciftci, O.N. In Vitro Bioaccessibility of Novel Low-Crystallinity Phytosterol Nanoparticles in Non-Fat and Regular-Fat Foods. *Food Res. Int.* 2019, 123, 27–35. [CrossRef]
- 78. Resende, D.; Costa Lima, S.A.; Reis, S. Nanoencapsulation Approaches for Oral Delivery of Vitamin A. *Colloids Surf. B Biointerfaces* **2020**, 193. [CrossRef]
- Xiang, C.; Gao, J.; Ye, H.; Ren, G.; Ma, X.; Xie, H.; Fang, S.; Lei, Q.; Fang, W. Development of Ovalbumin-Pectin Nanocomplexes for Vitamin D3 Encapsulation: Enhanced Storage Stability and Sustained Release in Simulated Gastrointestinal Digestion. *Food Hydrocoll.* 2020, 106. [CrossRef]
- 80. Gupta, C.; Arora, S.; Sharma, A.; Sharma, V. Evaluation of Effective Storage Conditions and In-Vitro Bioaccessibility of Vitamin A from Native and Modified Sodium Caseinate-Vitamin A Complexes. *LWT* **2019**, *111*, 284–290. [CrossRef]
- 81. Maurya, V.K.; Aggarwal, M. A Phase Inversion Based Nanoemulsion Fabrication Process to Encapsulate Vitamin D3 for Food Applications. *J. Steroid Biochem. Mol. Biol.* **2019**, *190*, 88–98. [CrossRef]
- Sahafi, S.M.; Goli, S.A.H.; Kadivar, M.; Varshosaz, J.; Shirvani, A. Pomegranate Seed Oil Nanoemulsion Enriched by α-Tocopherol; the Effect of Environmental Stresses and Long-Term Storage on Its Physicochemical Properties and Oxidation Stability. *Food Chem.* 2021, 345. [CrossRef]
- Hategekimana, J.; Chamba, M.V.M.; Shoemaker, C.F.; Majeed, H.; Zhong, F. Vitamin E Nanoemulsions by Emulsion Phase Inversion: Effect of Environmental Stress and Long-Term Storage on Stability and Degradation in Different Carrier Oil Types. *Colloids Surf. A Physicochem. Eng. Asp.* 2015, 483, 70–80. [CrossRef]
- Lv, S.; Gu, J.; Zhang, R.; Zhang, Y.; Tan, H.; McClements, D.J. Vitamin e Encapsulation in Plant-Based Nanoemulsions Fabricated Using Dual-Channel Microfluidization: Formation, Stability, and Bioaccessibility. J. Agric. Food Chem. 2018, 66, 10532–10542. [CrossRef]
- 85. Borba, C.M.; Tavares, M.N.; Macedo, L.P.; Araújo, G.S.; Furlong, E.B.; Dora, C.L.; Burkert, J.F.M. Physical and Chemical Stability of β-Carotene Nanoemulsions during Storage and Thermal Process. *Food Res. Int.* **2019**, *121*, 229–237. [CrossRef]
- 86. Sedaghat Doost, A.; Afghari, N.; Abbasi, H.; Nikbakht Nasrabadi, M.; Dewettinck, K.; van der Meeren, P. Nano-Lipid Carriers Stabilized by Hydrophobically Modified Starch or Sucrose Stearate for the Delivery of Lutein as a Nutraceutical Beverage Model. *Colloids Surf. A Physicochem. Eng. Asp.* **2020**, *605*. [CrossRef]
- Zimet, P.; Rosenberg, D.; Livney, Y.D. Re-Assembled Casein Micelles and Casein Nanoparticles as Nano-Vehicles for ω-3 Polyunsaturated Fatty Acids. *Food Hydrocoll.* 2011, 25, 1270–1276. [CrossRef]
- Zimet, P.; Livney, Y.D. Beta-Lactoglobulin and Its Nanocomplexes with Pectin as Vehicles for ω-3 Polyunsaturated Fatty Acids. *Food Hydrocoll.* 2009, 23, 1120–1126. [CrossRef]
- 89. Liu, M.; Yang, C.; Liu, E.; Zhang, F.; Meng, X.; Liu, B. Effect of Environmental Stresses on Physicochemical Properties of ALA Oil-in-Water Nanoemulsion System Prepared by Emulsion Phase Inversion. *Food Chem.* **2020**. [CrossRef]
- Rasti, B.; Jinap, S.; Mozafari, M.R.; Yazid, A.M. Comparative Study of the Oxidative and Physical Stability of Liposomal and Nanoliposomal Polyunsaturated Fatty Acids Prepared with Conventional and Mozafari Methods. *Food Chem.* 2012, 135, 2761–2770. [CrossRef] [PubMed]
- 91. Huang, J.; Wang, Q.; Li, T.; Xia, N.; Xia, Q. Nanostructured Lipid Carrier (NLC) as a Strategy for Encapsulation of Quercetin and Linseed Oil: Preparation and in Vitro Characterization Studies. *J. Food Eng.* **2017**, *215*, 1–12. [CrossRef]

- 92. de Sousa Lobato, K.B.; Paese, K.; Forgearini, J.C.; Guterres, S.S.; Jablonski, A.; de Oliveira Rios, A. Evaluation of Stability of Bixin in Nanocapsules in Model Systems of Photosensitization and Heating. *LWT Food Sci. Technol.* **2015**, *60*, 8–14. [CrossRef]
- Li, X.M.; Li, X.; Wu, Z.; Wang, Y.; Cheng, J.S.; Wang, T.; Zhang, B. Chitosan Hydrochloride/Carboxymethyl Starch Complex Nanogels Stabilized Pickering Emulsions for Oral Delivery of β-Carotene: Protection Effect and in Vitro Digestion Study. *Food Chem.* 2020, 315. [CrossRef]
- 94. Ma, L.; Zou, L.; McClements, D.J.; Liu, W. One-Step Preparation of High Internal Phase Emulsions Using Natural Edible Pickering Stabilizers: Gliadin Nanoparticles/Gum Arabic. *Food Hydrocoll.* **2020**, *100*. [CrossRef]
- Burgos-Díaz, C.; Opazo-Navarrete, M.; Soto-Añual, M.; Leal-Calderón, F.; Bustamante, M. Food-Grade Pickering Emulsion as a Novel Astaxanthin Encapsulation System for Making Powder-Based Products: Evaluation of Astaxanthin Stability during Processing, Storage, and Its Bioaccessibility. *Food Res. Int.* 2020, 134. [CrossRef] [PubMed]
- Giaconia, M.A.; dos Passos Ramos, S.; Pereira, C.F.; Lemes, A.C.; de Rosso, V.V.; Braga, A.R.C. Overcoming Restrictions of Bioactive Compounds Biological Effects in Food Using Nanometer-Sized Structures. *Food Hydrocoll.* 2020, 107, 105939.
- 97. Yao, M.; Xiao, H.; McClements, D.J. Delivery of Lipophilic Bioactives: Assembly, Disassembly, and Reassembly of Lipid Nanoparticles. *Annu. Rev. Food Sci. Technol.* **2014**, *5*, 53–81. [CrossRef]
- Bao, C.; Jiang, P.; Chai, J.; Jiang, Y.; Li, D.; Bao, W.; Liu, B.; Liu, B.; Norde, W.; Li, Y. The Delivery of Sensitive Food Bioactive Ingredients: Absorption Mechanisms, Influencing Factors, Encapsulation Techniques and Evaluation Models. *Food Res. Int.* 2019, 120, 130–140. [CrossRef]
- 99. Guri, A.; Gülseren, I.; Corredig, M. Utilization of Solid Lipid Nanoparticles for Enhanced Delivery of Curcumin in Cocultures of HT29-MTX and Caco-2 Cells. *Food Funct.* 2013, 4, 1410–1419. [CrossRef]
- Mitsou, E.; Dupin, A.; Sassi, A.H.; Monteil, J.; Sotiroudis, G.T.; Leal-Calderon, F.; Xenakis, A. Hydroxytyrosol Encapsulated in Biocompatible Water-in-Oil Microemulsions: How the Structure Affects in Vitro Absorption. *Colloids Surf. B Biointerfaces* 2019, 184. [CrossRef]
- 101. Gasa-Falcon, A.; Arranz, E.; Odriozola-Serrano, I.; Martín-Belloso, O.; Giblin, L. Delivery of β-Carotene to the in Vitro Intestinal Barrier Using Nanoemulsions with Lecithin or Sodium Caseinate as Emulsifiers. LWT 2021, 135. [CrossRef]
- Inapurapu, S.P.; Ibrahim, A.; Kona, S.R.; Pawar, S.C.; Bodiga, S.; Bodiga, V.L. Development and Characterization of ω-3 Fatty Acid Nanoemulsions with Improved Physicochemical Stability and Bioaccessibility. *Colloids Surf. A Physicochem. Eng. Asp.* 2020, 606. [CrossRef]
- 103. Edelman, R.; Engelberg, S.; Fahoum, L.; Meyron-Holtz, E.G.; Livney, Y.D. Potato Protein- Based Carriers for Enhancing Bioavailability of Astaxanthin. *Food Hydrocoll.* **2019**, *96*, 72–80. [CrossRef]
- 104. Rana, S.; Arora, S.; Gupta, C.; Kapila, S. Effect of Sodium Caseinate and Vitamin A Complexation on Bioaccessibility and Bioavailability of Vitamin A in Caco-2 Cells. *Food Res. Int.* **2019**, 121, 910–918. [CrossRef]
- Park, S.J.; Garcia, C. v.; Shin, G.H.; Kim, J.T. Development of Nanostructured Lipid Carriers for the Encapsulation and Controlled Release of Vitamin D3. *Food Chem.* 2017, 225, 213–219. [CrossRef] [PubMed]
- Ozturk, B.; Argin, S.; Ozilgen, M.; McClements, D.J. Nanoemulsion Delivery Systems for Oil-Soluble Vitamins: Influence of Carrier Oil Type on Lipid Digestion and Vitamin D3 Bioaccessibility. *Food Chem.* 2015, 187, 499–506. [CrossRef]
- 107. Yang, Y.; McClements, D.J. Vitamin E Bioaccessibility: Influence of Carrier Oil Type on Digestion and Release of Emulsified α-Tocopherol Acetate. *Food Chem.* **2013**, *141*, 473–481. [CrossRef]
- 108. Xu, F.; Pandya, J.K.; Chung, C.; McClements, D.J.; Kinchla, A.J. Emulsions as Delivery Systems for Gamma and Delta Tocotrienols: Formation, Properties and Simulated Gastrointestinal Fate. *Food Res. Int.* **2018**, *105*, 570–579. [CrossRef]
- Yuan, Y.; Li, H.; Liu, C.; Zhang, S.; Xu, Y.; Wang, D. Fabrication and Characterization of Lutein-Loaded Nanoparticles Based on Zein and Sophorolipid: Enhancement of Water Solubility, Stability, and Bioaccessibility. J. Agric. Food Chem. 2019. [CrossRef] [PubMed]
- Ha, T.V.A.; Kim, S.; Choi, Y.; Kwak, H.S.; Lee, S.J.; Wen, J.; Oey, I.; Ko, S. Antioxidant Activity and Bioaccessibility of Size-Different Nanoemulsions for Lycopene-Enriched Tomato Extract. *Food Chem.* 2015, 178, 115–121. [CrossRef]
- 111. Feng, S.; Wang, D.; Gan, L.; Shao, P.; Jiang, L.; Sun, P. Preparation and Characterization of Zein/Pectin-Based Phytosterol Nanodispersions and Kinetic Study of Phytosterol Release during Simulated Digestion in Vitro. *LWT* **2020**, *128*. [CrossRef]
- 112. Mcclements, D.J.; Xiao, H. Excipient Foods: Designing Food Matrices That Improve the Oral Bioavailability of Pharmaceuticals and Nutraceuticals. *Food Funct.* **2014**, *5*, 1320–1333. [CrossRef] [PubMed]
- 113. Salvia-Trujillo, L.; McClements, D.J. Enhancement of Lycopene Bioaccessibility from Tomato Juice Using Excipient Emulsions: Influence of Lipid Droplet Size. *Food Chem.* **2016**, *210*, 295–304. [CrossRef]
- 114. Yi, J.; Lam, T.I.; Yokoyama, W.; Cheng, L.W.; Zhong, F. Cellular Uptake of β-Carotene from Protein Stabilized Solid Lipid Nanoparticles Prepared by Homogenization-Evaporation Method. *J. Agric. Food Chem.* **2014**, *62*, 1096–1104. [CrossRef]
- 115. Dey, T. kumar; Ghosh, S.; Ghosh, M.; Koley, H.; Dhar, P. Comparative Study of Gastrointestinal Absorption of EPA & DHA Rich Fish Oil from Nano and Conventional Emulsion Formulation in Rats. *Food Res. Int.* **2012**, *49*, 72–79. [CrossRef]
- 116. Abuasal, B.S.; Lucas, C.; Peyton, B.; Alayoubi, A.; Nazzal, S.; Sylvester, P.W.; Kaddoumi, A. Enhancement of Intestinal Permeability Utilizing Solid Lipid Nanoparticles Increases γ-Tocotrienol Oral Bioavailability. *Lipids* **2012**, 47, 461–469. [CrossRef] [PubMed]

englanti	suomi
3D network	kolmiulotteinen verkosto (D= dimension, ulottuvuus)
absorption	imeytyminen, imeytyminen ruoansulatuskanavasta
accelerate; accelerated shelf-life	nopeutettu; nopeutettu säilyvyyskoe: menetelmä, jossa
stress test	säilvtysolosuhteita muuttamalla saadaan tuote vanhenemaan
	nopeammin
acrylamide	akryyliamidi
aerogel	aerogeeli: geelirakenne, jossa aeelin neste korvattu kaasulla
aftertaste	jälkimaku
aggregating: aggregation state	aggregoituminen: aggregaatiotila
alga, algae: algae oil	levä. levät: leväöliv
alpha-linolenic acid (ALA), $\alpha$ -	alfalinoleenihappo [rasvahappo 18:3(n=3)]
linolenic acid	
all-trans-retinol	all-trans-retinoli
animal study: animal model	eläinkoe: eläinmalli: eläinkoe, joka mallintaa ihmisen fysiologiga tai
	sairautta
anti-inflammatory	anti-inflammatorinen tulehdusta ehkäisevä
anticancer: anticarcinogen	svönää ehkäisevä: karsinogeenin (svönää aiheuttavan aineen)
anticalier, anticalenogen	vastavaikuttaja
antidiabetic (anti-diabetagenic)	diabetesta ehkäisevä
antimicrobial	antimikrohinen, mikrohien kasvua ehkäisevä
antioxidant	antinkrobinch, mikrobich kasva chkaiseva
annearance	
application	coveltaminen käyttö covelluckohde menetelmä
application	
aqueous environment	vesiyilipalisto, rulisaasti vetta sisaltava yilipalisto
	arami
arterioscierosis >> atheroscierosis	aleroskieroosi: valimoiden anlaulumiseen jonlava lauliprosessi,
accombly disaccombly	tässä uhtoon liittuminon hojoontuminon
assembly, disassembly	
attractive interaction	attraktio, vetovoima: positiivisesti ja negatiivisesti varautuneiden
attribute; attributed to	ominalsuus; selittya jollakin
betacarotene, β-carotene	beetakaroteeni (karotenoidi)
betalactoglobulin; β-lactoglobulin	beetalaktoglobuliini
betasitosterol, β-sitosterol	beetasitosteroli (kasvisteroli)
beverage	juoma
bilayer; lipid bilayer	kaksoiskerros; lipidikaksoiskerros
bile salt	sappisuola
bioaccessibility	biosaatavuus: se osuus ruoan sisältämästä aineesta, joka vapautuu
	ruoansulatuskanavaa jäljittelevissä olosuhteissa ja olisi elimistöön
	imeytyvässä muodossa
bioactive lipid	bioaktiivinen lipidi: lipidi, joka voi vaikuttaa ihmisen terveyteen
bioavailability; bioavailable	hyväksikäytettävyys: se osuus syödystä ravintoaineesta (tai muusta
	ravinnon sisältämästä aineesta), joka on elimistön solujen
	käytettävissä; hyväksikäytettävä
biopolymer	biopolymeeri: luonnossa esiintyvä orgaaninen polymeeri
bixin (a carotenoid)	biksiini (karotenoidi)
blood clotting, blood coagulation	veren hyytyminen

bloodstream	verenkierto
bridging	silloittuminen
bulk; bulk form, bulk oil	öljy öljymuodossa: korostettaessa sitä, ettei öljyä ole liuotettu tai
	dispergoitu mihinkään
by-pass	ohittaa
campesterol	kampesteroli (kasvisteroli)
capillary forces	kapillaarivoimat
capsule; nanocapsule	kapseli; nanokapseli
carboxymethyl starch	karboksimetyylitärkkelys
cardiovascular disease	sepelvaltimotauti
carotene; $\alpha$ -carotene, $\beta$ -carotene	karoteeni; alfakaroteeni; beetakaroteeni
carotenoid	karotenoidi
carrier; nanocarrier; carrier particle	kantaja, kantaja-aine; nanokantaja; kantajapartikkeli
casein; casein micelles, casein	kaseiini; kaseiinimiselli, kaseiininanopartikkeli
nanoparticles	
caseinate	kaseinaatti
cataract	harmaakaihi
cell line; Caco-2 cell line	solulinja; Caco-2 solulinja (imeytymistutkimuksissa käytettävä
	solumalli)
cell membrane	solukalvo
cell proliferation	solunjakautuminen
cellulose; nanocellulose	selluloosa; nanoselluloosa
charge	varaus
chemical; chemically	kemiallinen; kemiallisesti
chilling	jäähdytys
chitin; nanochitin	kitiini; nanokitiini
chitosan hydrochloride	kitosaanihydrokloridi
cholesterol-lowering	kolesterolia alentava, seerumin kolesterolipitoisuutta pienentävä
chylomicron	kylomikroni: lipoproteiinihiukkanen, joka kuljettaa suolesta
	imeytyneitä rasvoja elimistöön
coalesce; coalescence	yhtyä, yhdistyä; yhteensulautuminen, yhdistyminen
coating	päällystäminen, päällyste
cognitive function	kognitiivinen toiminto, tiedon käsittelyyn liittyvä toiminto
colloidal	kolloidinen: systeemi, jossa pieniä partikkeleita (kooltaan 1 nm – 1
	μm) jakautuneena (dispergoituneena) jatkuvaan faasiin
color stability	värin pysyvyys
compatibility	yhteensopivuus
complex; nanocomplex	kompleksi; nanokompleksi
composition	koostumus
concentric	samankeskinen: esim. ympyröitä/palloja/kehiä, joilla on yhteinen
	keskipiste tai akseli
condiment	maustekastike, mauste
conjugated linoleic acid (CLA)	konjugoitu linolihappo (konjugoidussa rasvahapossa kaksoissidokset
	ovat peräkkäisissä hiilissä)
consumer panelists	kuluttajapaneelin jäsenet: kuluttajapaneelin avulla mitataan
	esimerkiksi tuotteen hyväksyttävyyttä
consumption	kulutus, käyttö, syöminen ja juominen
conventional	tavanomainen, perinteinen
coronary heart disease	sepelvaltimotauti
creaming	kermoittuminen

crosslink (cross-link); crosslinked	ristisidos, siltasidos: sidoksia, jotka yhdistävät näytteen eri osia
(cross-linked)	<i>toisiinsa</i> ; silloitettu, ristisidottu
crude carotenoid extract	puhdistamaton karotenoidiuute
crystallize; crystalline structure	kiteytyä; kiderakenne
cytokine storm	sytokiinimyrsky: kudoksia tuhoava tulehdustila
d, diameter	läpimitta, halkaisija
dark leafy green vegetables	tummanvihreät lehtivihannekset
degenerative disease	rappeumasairaus
degrade; degradation	hajottaa, hajota; hajoaminen
delivery system	kuljetin: viittaa elintarvikkeeseen, jonka mukana haluttu
	komponentti syödään tai juodaan
depletion	loppuun kuluminen, tyhjentäminen
diet; dietary	ruokavalio; ruokavalioon liittyvä
digest; digestion; digestion model	sulattaa; ruoansulatus; ruoansulatusmalli
disperse, dispersed; dispersibility	dispergoitua, dispergoitunut: dispersiossa (kts. alla)
	dispergoituneena on se aine, joka on pieninä osasina jatkuvan faasin
	joukossa; dispergoituvuus
dispersion	dispersio: kahden toisiinsa liukenemattoman aineen seos, jossa yhtä
	ainetta on pieniksi osasiksi jakaantuneena toisessa aineessa, joka
	muodostaa jatkuvan faasin
docosahexaenoic acid (DHA)	dokosaheksaeenihappo [rasvahappo 22:6(n–3)]
domain	alue
drawback	haitta, huono puoli
edible	syötävä
efficacy	teho, vaikuttavuus
egg yolk	munankeltuainen
eicosapentaenoic acid (EPA)	eikosapentaeenihappo [rasvahappo 20:5(n-3)]
electrospinning	sähkökehräys
electrostatic	sähköstaattinen
emulsify; emulsification; emulsifier	emulgoida; emulgointi, emulsion valmistaminen; emulgointiaine,
	emulgaattori
emulsion	emulsio
encapsulate; encapsulated,	kapseloida; kapseloitu
enterocyte	enterosyytti, ohutsuolen epiteelisolu
enzyme	entsyymi
epithelial cell, epithelium cell	epiteelisolu, peittosolu: ihon ja limakalvojen pinnan solut
essential fatty acid	välttämätön rasvahappo: rasvahappo, jolla on välttämättömiä
	tehtäviä elimistössä, mutta jota elimistö ei itse syntetisoi, minkä
	takia se pitää saada ravinnosta
	Huom! Artikkelissa on virhe: CLA ei ole välttämätön rasvahappo.
	Varsinaiset valttamattomat rasvahapot ovat linolihappo ja
essential oil, flavor oil	eteerinen oljy, tuoksuoljy, aromioljy
esterity	esteroida, muodostaa esteri
evaporation	naintuminen, naindutus
excipient roods	ruoku, jolla el ilsessaan ploaktiivisuutta, mutta joka lisaa sen kanssa
ovnalling, ovnallad	syotyjen bloaktiivisten ynaistelaen nyvaksikäytettävyyttä
experiing; experied	erottammen; eroteitu
exterior	uikomen, uikopuolinen, uiko-
exilaci	Tuule, uuttaa

fabricated	valmistettu
fat soluble vitamin	rasvaliukoinen vitamiini
fiber; nanofiber	kuitu; nanokuitu
first-pass metabolism	ensikierron aineenvaihdunta: suolesta imeytyvän aineen muokkaus
	ennen sen pääsyä yleiseen verenkiertoon
flaxseed oil	pellavansiemenöljy
flocculation	flokkulaatio, kasaantuminen, ryppäiden muodostuminen
food application	elintarvikesovellus
food grade (food-grade)	elintarvikelaatuinen
food matrix (pl. matrices)	elintarvikkeen rakenne, elintarvikematriisi; ruoan sisältämien
	komponenttien muodostama kokonaisuus
fortify; fortification	täydentää, lisätä ravintoaineita; täydentäminen: ravintoaineen
	lisääminen elintarvikkeeseen tai sen raaka-aineeseen
fraction	fraktio, jae; erotus, erottelu, fraktiointi, jaottelu
free fatty acid (FFA)	vapaa rasvahappo, esteröitymätön rasvahappo
freeze drying	pakkaskuivaus
freeze-thawing	pakastus-sulatus
fucoxanthin	fukoksantiini (karotenoidi)
functional food	terveysvaikutteinen elintarvike
functionality; functional attributes	toiminnallisuus; toiminnalliset ominaisuudet
gammatocotrienol; γ-tocotrienol	gammatokotrienoli
gastric	maha-, mahalaukkuun liittyvä
gastrointestinal; gastrointestinal	ruoansulatuselimistöön liittyvä; ruoansulatuskanava,
tract (GIT); GIT model	mahasuolikanava; ruoansulatusmalli: <i>laboratoriomalli, joka</i>
	jäljittelee ruoansulatuksen tapahtumia
gel; nanogel	geeli; nanogeeli
gelatin	gelatiini, liivate
gliadin	gliadiini
glycosylation	glykosylaatio; tässä: sokereiden liittäminen proteiiniin
granola bar	myslipatukka
gravitational; gravitational	painovoimaan liittyvä; painovoiman aiheuttama erottelu tai
separation	erottuminen
gum arabic	arabikumi
gut	suoli
heat treatment	lämpökäsittely
high amylose corn starch	runsaasti amyloosia sisältävä maissitärkkelys
high-energy approach	paljon ulkoista energiaa vaativa valmistustapa
high-melting lecithin	korkeassa lämpötilassa sulava lesitiini
homogeneity; homogenization;	homogeenisuus: rakenteen tasaisuus; homogenointi;
homogenizer	homogenisaattori/homogenoija
human study	tässä: ihmisillä tehtävä kokeellinen tutkimus
hydrophilic / hydrophobic	hydrofiilinen, veteen suuntautuva / hydrofobinen, vesipakoinen, vettä hylkivä
hypolipidemic	hypolipideminen; tässä: seerumin kolesterolipitoisuuden
	pieneneminen (kohti tavoiteltavampaa pitoisuutta)
immune system;	immuunijärjestelmä; immuunivastetta muokkaava
immunomodulatory	
in vitro / in vivo	koeputkessa, laboratoriossa / elävässä elimistössä
infant formula	äidinmaidonkorvike

inflammation	tulehdus
ingestion	ravinnonotto, syöminen
ingredient	aines, ainesosa, elintarvikkeen valmistusaine
instability	epävakaus, epästabiilius
interaction	vuorovaikutus
interface; interfacial	rajapinta; rajapintaan liittyvä
intestinal fluid	suolineste
ionic strength	ionivahvuus
isomer; positional isomer;	isomeeri; paikkaisomeeri; geometrinen isomeeri
geometric isomer	
isoprenoid and isoprene side	isoprenoidi- ja isopreenisivuketjut
chains	
krill oil	krilliöljy (äyriäisöljy)
labile	labiili, epävakaa
liberate; liberation	vapauttaa; vapauttaminen
lining of the human GIT	tässä: suolen limakalvo
linoleic acid	linolihappo [rasvahappo 18:2(n–6)]
lipase	lipaasi (entsyymi)
lipid	lipidi
lipophilic	lipofiilinen, rasvahakuinen
liposome	liposomi
low-energy approach	vähän ulkoista energiaa vaativa valmistustapa
lupin	lupiini
lutein	luteiini (karotenoidi)
lycopene	lykopeeni (karotenoidi)
lymphatic system	lymfaattinen järjestelmä, imusuonisto
M-cell	M-solu: suolen immuunijärjestelmään liittyvä solu
matrix compatibility	matriisiyhteensopivuus
matrix (plural: matrices)	matriisi, rakenne
melting point	sulamispiste
metabolism; metabolic	aineenvaihdunta; aineenvaihduntaan liittyvä
metal ion	metalli-ioni
method	menetelmä
micelle; micellar	miselli; misellimäinen
microfluidization; microfluidizer	mikrofluidisointi: homogenointimenetelmä; mikrofluidisoija
micronutrient	mikroravintoaine: ihmiselle välttämättömät kivennäisaineet ja
	vitamiinit
microparticle	mikropartikkeli
milk analog	maidon tyyppinen, kasviperäinen juoma, esimerkiksi kaurajuoma
	("kauramaito")
mixed micelle	seosmiselli: rasvojen ruoansulatustuotteiden ja sappisuolojen
	muodostama yhteenliittyma, joka auttaa rasvojen
	ruoansulatustuottelaen pääsyä suolen ontelosta ravintoa imeyttäviin
melecule	
	molekyyii
(MG)	monoasyyngiyseron; monogiyseriai
	limaa tuottava (suolon) selu
nanoparticla	ninaa tuuttava (suulen) sulu
nanoparticle	nanopartikkeli
nanoporous	nanonuokoinen: systeemi, jossa nanokokoisia huokosia

nanostructured lipid carriers (NLCs)	nanorakenteiset lipidikantajat
nanotechnology	nanoteknologia
network	verkosto
neurological, nerval	neurologinen, hermostoon liittyvä
nutraceutical	Termille ei ole yleisesti hyväksyttyä määritelmää eikä suomenkielistä
	käännöstä. Termillä tarkoitetaan yleensä ruokia tai ruoasta peräisin
	olevia komponentteja, joilla on terveyttä edistäviä tai sairauksia
	ehkäiseviä tai hoitavia vaikutuksia.
off-flavor	virhemaku
oil droplet	öljypisara
oil-in-water	öljy vedessä
oil-soluble	rasvaliukoinen
omega-3 fatty acid	omega-3-rasvahappo
oral; orally administered	suu-, suuhun liittyvä; suun kautta annosteltu
ostheosynthetic	tässä: luunmuodostusta parantava
Ostwald ripening	Ostwaldin kypsyminen
ovalbumin	ovalbumiini: yksi kananmunan proteiineista
oxidation; oxidative stability	hapettuminen; hapettumiskestävyys: kestävyys hapettumista
	vastaan
oxygenated / nonoxygenated	happea sisältävä / happea sisältämätön
particle; particle size	partikkeli, hiukkanen; partikkelikoko
pasteurization	pastörointi
pectin	pektiini
phase; phase inversion	faasi; faasi-inversio: faasien vaihtuminen päinvastoin
phospholipid	fosfolipidi
physical; physically	fysikaalinen; fysikaalisesti
phytosterol	fytosteroli, kasvisteroli
Pickering emulsion	Pickering-emulsio: <i>emulsiotyyppi, jossa on kiinteitä partikkeleita</i>
plant storols and stanols	juusien rujupinnunu kasvistaralit ja, stapolit
plant based food	
plant-based 1000	nashipani nashiran nashiran asa malekuvliä
polar, non-polar, polar nead group	tässä kidemueden muutes teiseksi
polymorphic transition	
polyupsaturated fatty acid	polysäkkaitui monituuduttumätön rasuahanna
pore size	huokoskoko
preparation method	valmistusmenetelmä
pre-ovidant	
pro-vitamin A	
pros and cons	edut ja haitat
protrude	
nudding	vanukas
pulmonary	keuhkoien, keuhkoihin liittyvä
quercetin	kversetiini (flavonoidi/nolvfenoli)
guillaia saponin	kvillaian (suopapuun) saponiini
rat model	rottamalli (kts. animal model)
rate limiting	noneutta rajoittava
redispersed	uudelleen dispergoitu
refrigerator: refrigerated condition	jääkaappi: jääkaappiolosuhteet

reproduction	lisääntyminen (ihmisen tai eläimen)
repulsive	repulsoiva, hylkivä (vuorovaikutus)
respiratory tract disorder	hengityselimistön häiriötila, hengityselinsairaus
retention	pidättäminen, pidättyminen
retinol, all- <i>trans</i> retinol; retinyl	retinoli, all- <i>trans</i> -retinoli; retinyyliesteri
ester	
ripening	kypsyminen
ruminant	märehtijä
scale	mittakaava
seaweed	merilevä
sedimentation	sedimentaatio, saostuminen
sensory analysis	aistinvarainen arviointi: menetelmä, jossa arvioidaan näytteen
	ominaisuuksia aistien avulla (esimerkiksi katsellen, haistaen ja
	maistaen)
sensory acceptance, sensory	aistinvarainen hyväksyttävyys; aistinvaraisesti arvioitavat
acceptability; sensory attributes	ominaisuudet
shear; shearing	leikata, leikkaus-; leikkaus: tässä leikkauksella tarkoitetaan
	esimerkiksi sekoitusta
shelf-life	säilyvyysaika
small intestine	ohutsuoli
sodium caseinate	natriumkaseinaatti
solid lipid nanoparticle (SLN)	kiinteä lipidinanopartikkeli
solubilize; solubilization	liuottaa; liuotus
sonication	sonikointi, tässä: ultraäänikäsittely
sophorolipid	eräs glykolipidiluokka
spherical	pallomainen, pallon muotoinen
spherical spontaneous	pallomainen, pallon muotoinen spontaani: esim. spontaani reaktio (itsestään tapahtuva)
spherical spontaneous spray drying	pallomainen, pallon muotoinen spontaani: esim. spontaani reaktio (itsestään tapahtuva) sumutuskuivaus: menetelmä, jolla kuivataan nestemäisiä
spherical spontaneous spray drying	pallomainen, pallon muotoinen spontaani: esim. spontaani reaktio (itsestään tapahtuva) sumutuskuivaus: menetelmä, jolla kuivataan nestemäisiä elintarvikkeita (esimerkiksi maitojauheen valmistus)
spherical spontaneous spray drying stable; stability	pallomainen, pallon muotoinen spontaani: esim. spontaani reaktio (itsestään tapahtuva) sumutuskuivaus: menetelmä, jolla kuivataan nestemäisiä elintarvikkeita (esimerkiksi maitojauheen valmistus) stabiili, vakaa; stabiilius, vakaus
spherical spontaneous spray drying stable; stability starch	pallomainen, pallon muotoinen spontaani: esim. spontaani reaktio (itsestään tapahtuva) sumutuskuivaus: menetelmä, jolla kuivataan nestemäisiä elintarvikkeita (esimerkiksi maitojauheen valmistus) stabiili, vakaa; stabiilius, vakaus tärkkelys
spherical spontaneous spray drying stable; stability starch steric	pallomainen, pallon muotoinen spontaani: esim. spontaani reaktio (itsestään tapahtuva) sumutuskuivaus: menetelmä, jolla kuivataan nestemäisiä elintarvikkeita (esimerkiksi maitojauheen valmistus) stabiili, vakaa; stabiilius, vakaus tärkkelys steerinen
spherical spontaneous spray drying stable; stability starch steric sterilization	pallomainen, pallon muotoinen spontaani: esim. spontaani reaktio (itsestään tapahtuva) sumutuskuivaus: menetelmä, jolla kuivataan nestemäisiä elintarvikkeita (esimerkiksi maitojauheen valmistus) stabiili, vakaa; stabiilius, vakaus tärkkelys steerinen steerinen sterilointi: kuumennuskäsittely, jolla tuhotaan elintarvikkeessa
spherical spontaneous spray drying stable; stability starch steric sterilization	pallomainen, pallon muotoinen spontaani: esim. spontaani reaktio (itsestään tapahtuva) sumutuskuivaus: menetelmä, jolla kuivataan nestemäisiä elintarvikkeita (esimerkiksi maitojauheen valmistus) stabiili, vakaa; stabiilius, vakaus tärkkelys steerinen sterilointi: kuumennuskäsittely, jolla tuhotaan elintarvikkeessa olevat mikrobit ja bakteeri-itiöt
spherical spontaneous spray drying stable; stability starch steric sterilization steroid hormone	pallomainen, pallon muotoinen spontaani: esim. spontaani reaktio (itsestään tapahtuva) sumutuskuivaus: menetelmä, jolla kuivataan nestemäisiä elintarvikkeita (esimerkiksi maitojauheen valmistus) stabiili, vakaa; stabiilius, vakaus tärkkelys steerinen steerinen sterilointi: kuumennuskäsittely, jolla tuhotaan elintarvikkeessa olevat mikrobit ja bakteeri-itiöt steroidihormoni
spherical spontaneous spray drying stable; stability starch steric sterilization steroid hormone stigmasterol	pallomainen, pallon muotoinen spontaani: esim. spontaani reaktio (itsestään tapahtuva) sumutuskuivaus: menetelmä, jolla kuivataan nestemäisiä elintarvikkeita (esimerkiksi maitojauheen valmistus) stabiili, vakaa; stabiilius, vakaus tärkkelys steerinen steerinen sterilointi: kuumennuskäsittely, jolla tuhotaan elintarvikkeessa olevat mikrobit ja bakteeri-itiöt steroidihormoni stigmasteroli (kasvisteroli)
spherical spontaneous spray drying stable; stability starch steric sterilization steroid hormone stigmasterol stomach	pallomainen, pallon muotoinenspontaani: esim. spontaani reaktio (itsestään tapahtuva)sumutuskuivaus: menetelmä, jolla kuivataan nestemäisiä elintarvikkeita (esimerkiksi maitojauheen valmistus)stabiili, vakaa; stabiilius, vakaustärkkelyssteerinensterilointi: kuumennuskäsittely, jolla tuhotaan elintarvikkeessa olevat mikrobit ja bakteeri-itiötsteroidihormonistigmasteroli (kasvisteroli)mahalaukku
spherical spontaneous spray drying stable; stability starch steric sterilization steroid hormone stigmasterol stomach substitute	pallomainen, pallon muotoinenspontaani: esim. spontaani reaktio (itsestään tapahtuva)sumutuskuivaus: menetelmä, jolla kuivataan nestemäisiä elintarvikkeita (esimerkiksi maitojauheen valmistus)stabiili, vakaa; stabiilius, vakaustärkkelyssteerinensterilointi: kuumennuskäsittely, jolla tuhotaan elintarvikkeessa olevat mikrobit ja bakteeri-itiötsteroidihormonistigmasteroli (kasvisteroli)mahalaukku korvata, korvike
spherical spontaneous spray drying stable; stability starch steric sterilization steroid hormone stigmasterol stomach substitute surface area	pallomainen, pallon muotoinenspontaani: esim. spontaani reaktio (itsestään tapahtuva)sumutuskuivaus: menetelmä, jolla kuivataan nestemäisiä elintarvikkeita (esimerkiksi maitojauheen valmistus)stabiili, vakaa; stabiilius, vakaustärkkelyssteerinensterilointi: kuumennuskäsittely, jolla tuhotaan elintarvikkeessa olevat mikrobit ja bakteeri-itiötsteroidihormonistigmasteroli (kasvisteroli)mahalaukku korvata, korvikepinta-ala
spherical spontaneous spray drying stable; stability starch steric sterilization steroid hormone stigmasterol stomach substitute surface area surfactant; biosurfactant	pallomainen, pallon muotoinenspontaani: esim. spontaani reaktio (itsestään tapahtuva)sumutuskuivaus: menetelmä, jolla kuivataan nestemäisiä elintarvikkeita (esimerkiksi maitojauheen valmistus)stabiili, vakaa; stabiilius, vakaustärkkelyssteerinensterilointi: kuumennuskäsittely, jolla tuhotaan elintarvikkeessa olevat mikrobit ja bakteeri-itiötsteroidihormonistigmasteroli (kasvisteroli)mahalaukkukorvata, korvikepinta-alapinta-aktiivinen yhdiste; luonnosta peräisin oleva pinta-aktiivinen
spherical spontaneous spray drying stable; stability starch steric sterilization steroid hormone stigmasterol stomach substitute surface area surfactant; biosurfactant	pallomainen, pallon muotoinen spontaani: esim. spontaani reaktio (itsestään tapahtuva) sumutuskuivaus: menetelmä, jolla kuivataan nestemäisiä elintarvikkeita (esimerkiksi maitojauheen valmistus) stabiili, vakaa; stabiilius, vakaus tärkkelys steerinen steerinen sterilointi: kuumennuskäsittely, jolla tuhotaan elintarvikkeessa olevat mikrobit ja bakteeri-itiöt steroidihormoni stigmasteroli (kasvisteroli) mahalaukku korvata, korvike pinta-ala pinta-aktiivinen yhdiste; luonnosta peräisin oleva pinta-aktiivinen yhdiste
spherical spontaneous spray drying stable; stability starch steric sterilization steroid hormone stigmasterol stomach substitute surface area surfactant; biosurfactant suspend	pallomainen, pallon muotoinenspontaani: esim. spontaani reaktio (itsestään tapahtuva)sumutuskuivaus: menetelmä, jolla kuivataan nestemäisiä elintarvikkeita (esimerkiksi maitojauheen valmistus)stabiili, vakaa; stabiilius, vakaustärkkelyssteerinensterilointi: kuumennuskäsittely, jolla tuhotaan elintarvikkeessa olevat mikrobit ja bakteeri-itiötsteroidihormonistigmasteroli (kasvisteroli)mahalaukku korvata, korvikepinta-alapinta-aktiivinen yhdiste; luonnosta peräisin oleva pinta-aktiivinen yhdistesuspendoida, tehdä suspensio
spherical spontaneous spray drying stable; stability starch steric sterilization steroid hormone stigmasterol stomach substitute surface area surfactant; biosurfactant suspend tail	pallomainen, pallon muotoinen spontaani: esim. spontaani reaktio (itsestään tapahtuva) sumutuskuivaus: menetelmä, jolla kuivataan nestemäisiä elintarvikkeita (esimerkiksi maitojauheen valmistus) stabiili, vakaa; stabiilius, vakaus tärkkelys steerinen sterilointi: kuumennuskäsittely, jolla tuhotaan elintarvikkeessa olevat mikrobit ja bakteeri-itiöt steroidihormoni stigmasteroli (kasvisteroli) mahalaukku korvata, korvike pinta-ala pinta-aktiivinen yhdiste; luonnosta peräisin oleva pinta-aktiivinen yhdiste suspendoida, tehdä suspensio tässä: sivuketju
spherical spontaneous spray drying stable; stability starch steric sterilization steroid hormone stigmasterol stomach substitute surface area surfactant; biosurfactant suspend tail tailor; tailored	pallomainen, pallon muotoinen spontaani: esim. spontaani reaktio (itsestään tapahtuva) sumutuskuivaus: menetelmä, jolla kuivataan nestemäisiä elintarvikkeita (esimerkiksi maitojauheen valmistus) stabiili, vakaa; stabiilius, vakaus tärkkelys steerinen sterilointi: kuumennuskäsittely, jolla tuhotaan elintarvikkeessa olevat mikrobit ja bakteeri-itiöt steroidihormoni stigmasteroli (kasvisteroli) mahalaukku korvata, korvike pinta-ala pinta-aktiivinen yhdiste; luonnosta peräisin oleva pinta-aktiivinen yhdiste suspendoida, tehdä suspensio tässä: sivuketju räätälöidä, muokata; räätälöity
spherical spontaneous spray drying stable; stability starch steric steriic sterilization steroid hormone stigmasterol stomach substitute surface area surfactant; biosurfactant suspend tail tailor; tailored tetraterpenoid lipid	pallomainen, pallon muotoinenspontaani: esim. spontaani reaktio (itsestään tapahtuva)sumutuskuivaus: menetelmä, jolla kuivataan nestemäisiäelintarvikkeita (esimerkiksi maitojauheen valmistus)stabiili, vakaa; stabiilius, vakaustärkkelyssteerinensterilointi: kuumennuskäsittely, jolla tuhotaan elintarvikkeessa olevat mikrobit ja bakteeri-itiötsteroidihormonistigmasteroli (kasvisteroli)mahalaukku korvata, korvikepinta-alapinta-alapinta-aktiivinen yhdiste; luonnosta peräisin oleva pinta-aktiivinen yhdistesuspendoida, tehdä suspensiotässä: sivuketjuräätälöidä, muokata; räätälöity neljä terpeeniyksikköä sisältävä lipidi
spherical spontaneous spray drying stable; stability starch steric sterilization steroid hormone stigmasterol stomach substitute surface area surfactant; biosurfactant suspend tail tailor; tailored tetraterpenoid lipid texture; textural	pallomainen, pallon muotoinenspontaani: esim. spontaani reaktio (itsestään tapahtuva)sumutuskuivaus: menetelmä, jolla kuivataan nestemäisiäelintarvikkeita (esimerkiksi maitojauheen valmistus)stabiili, vakaa; stabiilius, vakaustärkkelyssteerinensterilointi: kuumennuskäsittely, jolla tuhotaan elintarvikkeessaolevat mikrobit ja bakteeri-itiötsteroidihormonistigmasteroli (kasvisteroli)mahalaukkukorvata, korvikepinta-alapinta-alasuspendoida, tehdä suspensiotässä: sivuketjuräätälöidä, muokata; räätälöityneljä terpeeniyksikköä sisältävä lipiditekstuuri, rakenne; rakenteeseen liittyvä aistittava tuntu
spherical spontaneous spray drying stable; stability starch steric steric sterilization steroid hormone stigmasterol stomach substitute surface area surface area surfactant; biosurfactant suspend tail tailor; tailored texture; textural thermal	pallomainen, pallon muotoinenspontaani: esim. spontaani reaktio (itsestään tapahtuva)sumutuskuivaus: menetelmä, jolla kuivataan nestemäisiäelintarvikkeita (esimerkiksi maitojauheen valmistus)stabiili, vakaa; stabiilius, vakaustärkkelyssteerinensterilointi: kuumennuskäsittely, jolla tuhotaan elintarvikkeessaolevat mikrobit ja bakteeri-itiötsteroidihormonistigmasteroli (kasvisteroli)mahalaukkukorvata, korvikepinta-alapinta-alapinta-aktiivinen yhdiste; luonnosta peräisin oleva pinta-aktiivinenyhdistesuspendoida, tehdä suspensiotässä: sivuketjuräätälöidä, muokata; räätälöityneljä terpeeniyksikköä sisältävä lipiditekstuuri, rakenne; rakenteeseen liittyvä aistittava tuntutermaalinen, lämpöön liittyvä, lämpö-
spherical spontaneous spray drying stable; stability starch steric sterilization steroid hormone stigmasterol stomach substitute surface area surfactant; biosurfactant suspend tail tailor; tailored tetraterpenoid lipid texture; textural thermal thermal	pallomainen, pallon muotoinen spontaani: esim. spontaani reaktio (itsestään tapahtuva) sumutuskuivaus: menetelmä, jolla kuivataan nestemäisiä elintarvikkeita (esimerkiksi maitojauheen valmistus) stabiili, vakaa; stabiilius, vakaus tärkkelys steerinen sterilointi: kuumennuskäsittely, jolla tuhotaan elintarvikkeessa olevat mikrobit ja bakteeri-itiöt steroidihormoni stigmasteroli (kasvisteroli) mahalaukku korvata, korvike pinta-ala pinta-aktiivinen yhdiste; luonnosta peräisin oleva pinta-aktiivinen yhdiste suspendoida, tehdä suspensio tässä: sivuketju räätälöidä, muokata; räätälöity neljä terpeeniyksikköä sisältävä lipidi tekstuuri, rakenne; rakenteeseen liittyvä aistittava tuntu termaalinen, lämpöön liittyvä, lämpö- termodynaaminen
spherical spontaneous spray drying stable; stability starch steric sterilization steroid hormone stigmasterol stomach substitute surface area surfactant; biosurfactant suspend tail tailor; tailored tetraterpenoid lipid texture; textural thermal thermodynamical tocopherol	pallomainen, pallon muotoinenspontaani: esim. spontaani reaktio (itsestään tapahtuva)sumutuskuivaus: menetelmä, jolla kuivataan nestemäisiäelintarvikkeita (esimerkiksi maitojauheen valmistus)stabiili, vakaa; stabiilius, vakaustärkkelyssteerinensterilointi: kuumennuskäsittely, jolla tuhotaan elintarvikkeessaolevat mikrobit ja bakteeri-itiötsteroidihormonistigmasteroli (kasvisteroli)mahalaukkukorvata, korvikepinta-alapinta-aktiivinen yhdiste; luonnosta peräisin oleva pinta-aktiivinenyhdistesuspendoida, tehdä suspensiotässä: sivuketjuräätälöidä, muokata; räätälöityneljä terpeeniyksikköä sisältävä lipiditekstuuri, rakenne; rakenteeseen liittyvä aistittava tuntutermodynaaminentokoferoli

trap; trapped	sulkea sisään; suljettu jonkin sisään
triglyceride (TG); long-chain	triglyseridi (triasyyliglyseroli); pitkäketjuisia rasvahappoja sisältävä
triglyceride (LCT); medium-chain	triglyseridi; keskipitkäketjuisia rasvahappoja sisältävä triglyseridi
triglyceride (MCT)	
unstable	epästabiili, epävakaa
van der Waals	van der Waals; <i>sidostyyppi</i>
viable	tässä: toimintakykyinen
viscosity	viskositeetti, nesteen jähmeys
visual	visuaalinen, tässä: näkökykyyn liittyvä
vitamin A	A-vitamiini
vitamin D; vitamin D <sub>2</sub>	D-vitamiini; D <sub>2</sub> -vitamiini (ergokalsiferoli); D <sub>3</sub> -vitamiini
(ergocalciferol); vitamin D₃	(kolekalsiferoli); 1,25-dihydroksi-D₃-vitamiini
(cholecalciferol); 1,25-	
dihydroxyvitamin D <sub>3</sub>	
vitamin E	E-vitamiini
vitamin K (phylloquinones and	K-vitamiini (fyllokinonit ja menakinonit)
menaquinones)	
water solubility	vesiliukoisuus
wet food products	kosteat tai nestemäiset elintarvikkeet
whey	hera
xanthophyll	ksantofylli (happea sisältävä karotenoidi)
zeaxanthin	zeaksantiini (karotenoidi)
zein	tseiini: eräs maissin proteiini



β-sitosterol



Stigmasterol



Campesterol



cis-9, trans-11 conjugated linoleic acid



trans-10, cis-12 conjugated linoleic acid